be optimized to save the most valves with the expected long-term durability. The authors clearly have developed a laudable technique that effectively results in durable valves. This is due in part to a respect for healthy-appearing native cusp morphology and humility to avoid excessive manipulation. In the context of proper valve selection and otherwise normal leaflets, cusp repair may

indeed reinforce the historical surgical tenant that perfect is the enemy of good.

Reference

1. Lau C, Wingo M, Rahouma M, Ivascu N, Iannacone E, Kamel M, et al. Valve-sparing root replacement in patients with bicuspid aortopathy: an analysis of cusp repair strategy and valve durability. *J Thorac Cardiovasc Surg*. 2021;161:469-78.

See Article page 469.

Commentary: To repair or not repair—that is the question

Kanika Kalra, MD, and Edward P. Chen, MD

Kanika Kalra, MD, and Edward P. Chen, MD

Aortic valve competency is dependent on multiple anatomic components involving the aortic root and valve cusps. Aortic insufficiency (AI) in bicuspid aortic valve (BAV) anatomy and aortic root pathology occurs as a result of abnormal cusp configuration as well as abnormal aortic root geometry.¹ Performing valve-sparing root replacement (VSRR) in patients with BAV AI allows correction of these components while establishing stable root geometry. Any moderate to severe AI after valve resuspension warrants correction, however, the question arises, whether mild AI should be aggressively repaired.

Lau and colleagues,² in this issue of *The Journal of Thoracic and Cardiovascular Surgery*, report their experience in 66 BAV patients who underwent VSRR, with or without aortic valve repair. The Cornell group has adopted a conservative approach to cusp repair and examined the outcomes of 51 BAV patients who did not receive cusp repair during VSRR compared with 15 patients who received cusp repair. In this series, most of the patients in the nonrepair group had 0 to mild AI (43/51), whereas most of the repair group had moderate or greater AI (11/15). Outcomes are outstanding and there was no

CENTRAL MESSAGE

Cusp repair is successfully done in insufficient bicuspid aortic valves during valve-sparing root replacement. Whether a conservative approach of noncusp repair is superior warrants further study.

operative mortality or major neurologic, renal, or cardiopulmonary complications.

At 3.5-year follow-up, echocardiography did not show any statistically significant differences in AI or aortic stenosis (AS) between the groups. Of the 51 patients who did not undergo cusp repair, 44 patients had trivial to mild AI at follow-up, indicating there was minimal progression of AI when left uncorrected. In 15 patients who received cusp repair, 13 patients had mild or less AI at the latest follow-up, indicating a significant improvement in valve function. Altogether, the authors conclude that cusp repair can be performed with adequate durability and safety in the setting of AI, and their outcomes might be equivalent in such patients compared with those who did not receive any valve repair.

From the Division of Cardiothoracic Surgery, Department of Surgery, Emory University School of Medicine, Atlanta, Ga.

Disclosures: Authors have nothing to disclose with regard to commercial support.

Received for publication Oct 28, 2019; accepted for publication Oct 29, 2019; available ahead of print Nov 22, 2020.

Address for reprints: Edward P. Chen, MD, Division of Cardiothoracic Surgery, Emory University, Atlanta, GA 30322 (E-mail: edward.p.chen@emory.edu).

J Thorac Cardiovasc Surg 2021;161:480-1

0022-5223/\$36.00

Copyright © 2019 by The American Association for Thoracic Surgery

<https://doi.org/10.1016/j.jtcvs.2019.10.189>

At our institution, we have adopted a more aggressive approach to repair all valves with AI with freedom from AI at 5 years being 93% and freedom for aortic valve replacement being 94%.³ Bavaria and colleagues also reported a similar aggressive approach, and reported very minimal AI recurrence at midterm follow-up.⁴ Our experience along with several others⁵ confirm the long-term safety of cusp repair, with acceptable rates of failure and reoperation.

The current study also confirms similar outcomes after cusp repair, albeit in a small group of patients at midterm follow-up. Despite these excellent outcomes with cusp repair, surprisingly the Kaplan-Meier curves indicate that 35% of patients in the cusp repair group had some AI immediately after surgery, and freedom from AI in the cusp repair group worsened with time. These results are potentially misleading because of the small number of cusp repair patients. Thus, the question of whether aggressive cusp repair versus a more conservative approach is superior in the long term may need further investigation. Experimental studies on the effect of cusp repair on mechanics of the valve and its degenerative potential may be contributory.

In summary, the Cornell aortic team should be commended for taking a highly unique and individualized approach to VSRR for correction of AI in BAV anatomy. This group has shown that VSRR for BAV AI can be done using a conservative approach with commendable midterm results.

References

1. Ram D, Bouhout I, Karliova I, Schneider U, El-Hamamsy I, Schäfers HJ. Concepts of bicuspid aortic valve repair - a review. *Ann Thorac Surg*. October 11, 2019 [Epub ahead of print].
2. Lau C, Wingo M, Rahouma M, Ivancu N, Iannacone E, Kamel M, et al. Valve-sparing root replacement in patients with bicuspid aortopathy: an analysis of cusp repair strategy and valve durability. *J Thorac Cardiovasc Surg*. 2021; 161:469-78.
3. Kayatta MO, Leshnower BG, McPherson L, Binongo JN, Lasanajak Y, Chen EP. Valve-sparing root replacement provides excellent midterm outcomes for bicuspid valve aortopathy. *Ann Thorac Surg*. 2019;107:499-504.
4. Bavaria JE, Desai N, Szeto WY, Komlo C, Rhode T, Wallen T, et al. Valve-sparing root reimplantation and leaflet repair in a bicuspid aortic valve: comparison with the 3-cusp David procedure. *J Thorac Cardiovasc Surg*. 2015;149:S22-8.
5. Schneider U, Feldner SK, Hofmann C, Schöpe J, Wagenpfeil S, Giebels C, et al. Two decades of experience with root remodeling and valve repair for bicuspid aortic valves. *J Thorac Cardiovasc Surg*. 2017;153:S65-71.

See Article page 469.

Commentary: Valve-sparing root replacement in bicuspid valves—more than technique

Dawn S. Hui, MD,^a and Kim I. de la Cruz, MD^b

A number of giants in the field of cardiothoracic surgery have, in our lifetime, advanced and elevated the discipline of aortic root surgery, with the ultimate goal of preserving native valve tissue and durable long-term function.¹ The knowledge gained and the techniques developed by these pioneers more recently have been applied to valve-sparing

Dawn S. Hui, MD, and Kim I. de la Cruz, MD, FACS, FCCP

CENTRAL MESSAGE

The optimal approach to cusp repair in valve-sparing aortic root operations for bicuspid aortic valve patients remains an area of investigation.

root (VSR) operations in patients with bicuspid aortic valves (BAV) and root aortopathy. In deciding which patients with BAV are candidates for VSR, one axiom on

From the ^aDepartment of Cardiothoracic Surgery, University of Texas Health Science Center at San Antonio; and ^bMethodist Heart Hospital San Antonio, San Antonio, Tex.

Disclosures: Authors have nothing to disclose with regard to commercial support. Received for publication Dec 6, 2019; revisions received Dec 6, 2019; accepted for publication Dec 7, 2019; available ahead of print Jan 3, 2020.

Address for reprints: Dawn S. Hui, MD, 7703 Floyd Curl Dr, Suite 211L, San Antonio, TX 78229 (E-mail: huid@uthscsa.edu).

J Thorac Cardiovasc Surg 2021;161:481-2
0022-5223/\$36.00

Copyright © 2019 by The American Association for Thoracic Surgery
<https://doi.org/10.1016/j.jtcvs.2019.12.033>