used in parts of Asia, and we certainly do not have all the answers there either.

References

1. Deckers JW, van Domburg RT, Akkerhuis M, Nauta ST. Relation of admission glucose levels, short- and long-term (20-year) mortality after acute myocardial infarction. *Am J Cardiol*. 2013;112:1306-10.
2. Malmberg K, Rydén L, Wedel H, Birkeland K, Bootsma A, Dirksen K, et al; DIGAMI 2 Investigators. Intense metabolic control by means of insulin in patients with diabetes mellitus and acute myocardial infarction (DIGAMI 2): effects on mortality and morbidity. *Eur Heart J*. 2005;26:650-61.
3. Chu LM, Osipov RM, Robich MP, Feng J, Oyamada S, Bianchi C, et al. Is hyperglycemia bad for the heart during acute ischemia? *J Thorac Cardiovasc Surg*. 2010;140:1345-52.
4. Charles EJ, Tian Y, Zhang A, Wu D, Mehaffey JH, Gigliotti JC, et al. Pulsed ultrasound attenuates the hyperglycemic exacerbation of myocardial ischemia-reperfusion injury. *J Thorac Cardiovasc Surg*. 2021;161:e297-306.
5. Gigliotti JC, Huang L, Bajwa A, Ye H, Mace EH, Hossack JA, et al. Ultrasound modulates the splenic neuroimmune axis in attenuating AKI. *J Am Soc Nephrol*. 2015;26:2470-81.

See Article page e297.

Commentary: The new kid in town: Pulsed ultrasound to prevent ischemia–reperfusion injury

Friedhelm Beyersdorf, MD

Ischemia–reperfusion injury (IRI) is a major clinical problem in all fields of medicine, especially in cardiac surgery and cardiology. Major improvements in understanding the pathophysiology, the causes, and the consequences have been achieved over the last decades.¹⁻³ Charles and colleagues⁴ from Irving Kron's group in Charlottesville are to be congratulated for drawing our attention to a new nonpharmacologic, noninvasive therapy option to reduce infarct size during acute myocardial infarction (MI). They provide promising data that pulsed ultrasound (pUS) is capable of attenuating the hyperglycemic exacerbation of myocardial IRI via the cholinergic anti-inflammatory pathway.⁴

Acute hyperglycemia in patients with MI (1) is independently associated with greater mortality and larger infarct size⁵ and (2) leads to the stimulation of splenic leukocytes through activation of the nicotinamide adenine dinucleotide

Friedhelm Beyersdorf, MD

CENTRAL MESSAGE

Pulsed ultrasound is a promising new approach to limit ischemia–reperfusion-injury.

phosphate oxidase pathway, which plays an important role in leukocyte-mediated reperfusion injury.⁶ Interestingly, reports show that pUS modulates splenic leukocytes into an anti-inflammatory phenotype.⁷

The strength of the study is that the authors not only report a significant reduction in infarct size in mice with acute hyperglycemia and 20 minutes of regional ischemia followed by 60 minutes of reperfusion, but that they provide insight into the molecular mechanisms of these beneficial effects. Splenic and neck pUS seem to reduce the splenic inflammatory response associated with hyperglycemic exacerbation of MI through vagus and acetylcholine signalling.⁴

The authors also point out 2 additional significant elements: (1) if neck pUS was applied 24 hours before IRI, it was as protective as if applied just before coronary occlusion; (2) when neck pUS was performed 10 minutes after reperfusion had started, no protective effect could be seen. In other words, the positive effects of modulation of splenic

From the Department of Cardiovascular Surgery, University Heart Center Freiburg–Bad Krozingen, and Albert-Ludwigs-University Freiburg, Freiburg, Germany.

Disclosures: Dr Beyersdorf is founder and shareholder of the start-up company “Resuscitec” of the University of Freiburg, Germany.

Received for publication Nov 4, 2019; revisions received Nov 4, 2019; accepted for publication Nov 4, 2019; available ahead of print Nov 27, 2019.

Address for reprints: Friedhelm Beyersdorf, MD, Department of Cardiovascular Surgery, University Heart Center Freiburg–Bad Krozingen, Hugstetterstr. 55, D-79106 Freiburg, Germany (E-mail: friedhelm.beyersdorf@uniklinik-freiburg.de).

J Thorac Cardiovasc Surg 2021;161:e308-9

0022-5223/\$36.00

Copyright © 2019 by The American Association for Thoracic Surgery

<https://doi.org/10.1016/j.jtcvs.2019.11.019>

leukocytes via neck pUS into an anti-inflammatory phenotype lasts at least 24 hours. However, once reperfusion injury had occurred (for 10 minutes), neck pUS could not reverse the reperfusion injury. Reversal of reperfusion injury—the holy grail of research in IRI—is not possible even with this new approach to prevent IRI. The authors are to be commended on adding this important information to their study.

The limitations of the study are obvious, and the authors have summarized most of them in the Discussion. The short reperfusion phase of 20 minutes and no further follow-up prevents drawing conclusions in the clinical setting. Potential systemic effects of neck and splenic pUS could not be assessed in the current experimental small animal model.

Nevertheless, the authors are to be congratulated for exploring a new path in preventing IRI and adding basic research to shed some light on the pathophysiologic mechanisms to this nonintuitive, innovative approach. The reduced interest of surgeons in basic science and the detrimental effects of this development for the future of surgery has been clearly expressed in a recent editorial in *Nature*.⁸ A transatlantic editorial about the same topic has described the reasons and possible solutions for this dilemma.⁹ This paper

by Charles and colleagues⁴ is a very good example of merging research on clinically relevant problems with basic science tools.

References

1. Buckberg GD. Studies of controlled reperfusion. I. When is cardiac muscle damaged irreversibly? *J Thorac Cardiovasc Surg*. 1986;92:483-7.
2. Taunyane IC, Benk C, Beyersdorf F, Foerster K, Schmitz HC, Wittmann K, et al. Preserved brain morphology after controlled automated reperfusion of the whole body following normothermic circulatory arrest time of up to 20 minutes. *Eur J Cardiothorac Surg*. 2016;50:1025-34.
3. Wu MY, Yian GT, Liao WT, Tsai APY, Cheng YL, Cheng PW, et al. Current mechanistic concepts in ischemia and reperfusion injury. *Cell Physiol Biochem*. 2018; 46:1650-67.
4. Charles EJ, Tian Y, Zhang A, Wu D, Mehaffey JH, Gigliotti JC, et al. Pulsed ultrasound attenuates the hyperglycemic exacerbation of myocardial ischemia-reperfusion injury. *J Thorac Cardiovasc Surg*. 2021;161:e297-306.
5. Deckers JW, van Domburg RT, Akkerhuis M, Nauta ST. Relation of admission glucose levels, short- and long-term (20-year) mortality after acute myocardial infarction. *Am J Cardiol*. 2013;112:1306-10.
6. Tian Y, French BA, Kron IL, Yang Z. Splenic leukocytes mediate the hyperglycemic exacerbation of myocardial infarct size in mice. *Bas Res Cardiol*. 2015;110:39.
7. Gigliotti JC, Huang L, Bajwa A, Ye H, Mace EH, Hossack JA, et al. Ultrasound modulates the splenic neuroimmune axis in attenuating AKI. *J Am Soc Nephrol*. 2015;26:2470-81.
8. More surgeons must start doing basic science. *Nature*. 2017;544:393-4.
9. Ikonomidis JS, Menasché P, Kreisel D, Sellke FW, Woo YJ, Colson YL. Attrition of the cardiothoracic surgeon-scientist: definition of the problem and remedial strategies. *J Thorac Cardiovasc Surg*. 2019;158:504-8.