6. Blommaert D, Gonzalez M, Mucumbitsi J, Gurné O, Evrard P, Buche M, et al. Effective prevention of atrial fibrillation by continuous atrial overdrive pacing after coronary artery bypass surgery. *J Am Coll Cardiol*. 2000;35:1411-5.
7. Greenberg MD, Katz NM, Iuliano S, Tempesta BJ, Solomon AJ. Atrial pacing for the prevention of atrial fibrillation after cardiovascular surgery. *J Am Coll Cardiol*. 2000;35:1416-22.
8. Budeus M, Feindt P, Gams E, Wieneke H, Sack S, Erbel R, et al. Beta-blocker prophylaxis for atrial fibrillation after coronary artery bypass grafting in patients with sympathovagal imbalance. *Ann Thorac Surg*. 2007;84:61-6.
9. Elgendy IY, Mahmoud A, Huo T, Beaver TM, Bavry AA. Meta-analysis of 12 trials evaluating the effects of statins on decreasing atrial fibrillation after coronary artery bypass grafting. *Am J Cardiol*. 2015;115:1523-8.
10. Mehaffey JH, Hawkins RB, Byler M, Smith J, Kern JA, Kron I, et al. Amiodarone protocol provides cost-effective reduction in postoperative atrial fibrillation. *Ann Thorac Surg*. 2018;105:1697-702.

See Article page 1803.

Commentary: Postoperative atrial fibrillation: “No magic bullet”

Scott M. Goldman, MD

In this issue of the *Journal*, Almassi and colleagues¹ reference a 2012 article² analyzing a subgroup of the Veterans Affairs Randomized On/Off Bypass follow-up study (ROOBY-FS).³ In 2012, Almassi’s group compared patients who developed atrial fibrillation during the hospitalization for the index procedure—postoperative atrial fibrillation (POAF)—and those who did not. They had previously reported risk-adjusted poorer outcomes and increase cost at a 1-year follow-up in patients who developed POAF.⁴ When analyzed at 5 years, these differences were not evident. There were no differences in the incidence of POAF between on-pump and off-pump coronary artery bypass. Beyond the first year, there were no differences in cost between those who developed POAF and those who did not. Almost all the increased cost at 1 year was attributed to those costs associated with the occurrence of POAF during the original hospitalization. As noted in the unadjusted data, the POAF group was older and had more complex comorbidities. This increased risk has also been demonstrated in other studies.⁵

By risk-adjusting the data, this study sheds more light on the patient population that is at increased risk for the development of POAF. They are older and have more

Scott M. Goldman, MD

CENTRAL MESSAGE

Only by considering the risk of postoperative atrial fibrillation in each patient and pursuing a holistic strategy are we likely to make a dent in the 30% incidence rate.

comorbidities, and it is evident from this and other studies^{5,6} that the burden surrounding POAF occurs during index hospitalization and the early postdischarge period.

Over the last 3 decades, the incidence of POAF has remained near 30% despite overall improvements in outcomes and postoperative care.⁶ There have been many attempts to identify the single intervention that will have a major impact on this. These include both pharmacologic and operative interventions.

Operative modifications that have been tried include posterior pericardiotomy, left atrial appendage exclusion, and anterior fat pad preservation.⁷⁻⁹ These have not proven effective. Left atrial appendage exclusion may increase the rate of POAF.^{6,10} Atrial and biatrial pacing have been studied, with mixed results.¹¹

The administration of beta blockers is the only class 1a indicated pharmacologic intervention for the prevention

From the Lankenau Heart Institute, Lankenau Medical Center, Wynnewood, Pa.
Disclosures: Author has nothing to disclose with regard to commercial support.
Received for publication Nov 13, 2019; revisions received Nov 13, 2019; accepted for publication Nov 13, 2019; available ahead of print Dec 7, 2019.

Address for reprints: Scott M. Goldman, MD, Lankenau Heart Institute, Lankenau Medical Center, 100 Lancaster Ave, Wynnewood, PA 19096 (E-mail: goldmans@mlhs.org).

J Thorac Cardiovasc Surg 2021;161:1812-3

0022-5223/\$36.00

Copyright © 2019 by The American Association for Thoracic Surgery

<https://doi.org/10.1016/j.jtcvs.2019.11.067>

of POAF with 82% of patients undergoing cardiac surgery receiving them.¹² The use of amiodarone for prevention of POAF has been shown to be effective in a large review of multiple clinical studies.¹³

It is clear that the burden of POAF is concentrated in the recovery period, and that is where our efforts should be focused. Is there any hope of changing this constant rate of POAF? Looking for the magic bullet seems to have failed. Perhaps a programmatic quality approach to this problem should be considered. First, we need to identify the patients at higher risk for the development of POAF, as identified in this and other studies. These include older patients, patients with more comorbidities, and patients requiring more complex procedure. The risk of POAF and its morbidity and high cost should be considered when deciding between surgery and less invasive transcatheter therapies. Converting complex procedures to less invasive hybrid strategies may be considered. These include hybrid coronary artery bypass and hybrid minimally invasive valve surgeries. Adding amiodarone to the pharmacologic regimen is likely to be effective when surgery is the best option for these patients at high risk for POAF.

Only by considering the risk of POAF in each patient and perusing a holistic strategy are we likely to make a dent in the 30% incidence rate, and as stated in the article, “future quality improvement strategies should focus primarily upon the preoperative and perioperative strategies to lessen the burden of POAF, and post-discharge care of POAF patients within the first post-CABG year period to potentially improve early postoperative clinical outcomes and costs.”

References

1. Almassi GH, Hawkins RB, Bishawi M, Shroyer AL, Hattler B, Quin JA, et al. New-onset postoperative atrial fibrillation impact on 5-year clinical outcomes and costs. *J Thorac Cardiovasc Surg.* 2021;161:1803-10.e3.
2. Almassi GH, Pecs SA, Collins JF, Shroyer AL, Zenati MA, Grover FL. Predictors and impact of postoperative atrial fibrillation on patients' outcomes: a report from the Randomized On Versus Off Bypass trial. *J Thorac Cardiovasc Surg.* 2012;143:93-102.
3. Shroyer AL, Hattler B, Wagner TH, Collins JF, Baltz JH, Quin JA, et al. Five-year outcomes after on-pump and off-pump coronary-artery bypass. *N Engl J Med.* 2017;377:623-32.
4. Almassi GH, Wagner TH, Carr B, Hattler B, Collins JF, Quin JA, et al. Postoperative atrial fibrillation impacts on costs and one-year clinical outcomes: the Veterans Affairs Randomized On/Off Bypass Trial. *Ann Thorac Surg.* 2015;99:109-14.
5. Akintoye E, Sellke F, Marchioli R, Tavazzi L, Mozaffarian D. Factors associated with postoperative atrial fibrillation and other adverse events after cardiac surgery. *J Thorac Cardiovasc Surg.* 2018;155:242-51.e10.
6. Greenberg JW, Lancaster TS, Schuessler RB, Melby SJ. Postoperative atrial fibrillation following cardiac surgery: a persistent complication. *Eur J Cardiothorac Surg.* 2017;52:665-72.
7. Melduni RM, Schaff HV, Lee HC, Gersh BJ, Noseworthy PA, Bailey KR, et al. Impact of left atrial appendage closure during cardiac surgery on the occurrence of early postoperative atrial fibrillation, stroke, and mortality: a propensity score-matched analysis of 10633 patients. *Circulation.* 2017;135:366-78.
8. Kaygin MA, Dag O, Güneş M, Senocak M, Limandal HK, Aslan U, et al. Posterior pericardiotomy reduces the incidence of atrial fibrillation, pericardial effusion, and length of stay in hospital after coronary artery bypasses surgery. *Tohoku J Exp Med.* 2011;225:103-8.
9. Bisleri G, Bottio T, Morgan JA, Muneretto C. Preservation of the anterior fat pad and incidence of postoperative atrial fibrillation following coronary surgery. *J Am Coll Cardiol.* 2005;45:1308.
10. Kongmalai P, Karunasumetta C, Kuptarnond C, Prathanee S, Taksinachanej S, Intanoo W, et al. The posterior pericardiotomy. Does it reduce the incidence of postoperative atrial fibrillation after coronary artery bypass grafting? *J Med Assoc Thai.* 2014;97(Suppl 10):S97-104.
11. Gerstenfeld EP, Hill MR, French SN, Mehra R, Rofino K, Vander Salm TJ, et al. Evaluation of right atrial and biatrial temporary pacing for the prevention of atrial fibrillation after coronary artery bypass surgery. *J Am Coll Cardiol.* 1999;33:1981-8.
12. Hillis LD, Smith PK, Anderson JL, Bittl JA, Bridges CR, Byrne JG, et al. 2011 ACCF/AHA guideline for coronary artery bypass graft surgery: executive summary: a report of the American College of Cardiology Foundation/American Heart Association task force on practice guidelines. *Circulation.* 2011;124:2610-42.
13. Arsenault KA, Yusuf AM, Crystal E, Healey JS, Morillo CA, Nair GM, et al. Interventions for preventing post-operative atrial fibrillation in patients undergoing heart surgery. *Cochrane Database Syst Rev.* 2013;1:CD003611.