The predictive value of the recently proposed SAFINOUS score still needs to be validated in a large multicenter cohort to assess the performance of the tool, particularly in the extremes of the risk spectrum. As with any risk score, it is virtually impossible to take all potential risk factors into consideration. For clinical decision making and surgery planning of CABG patients, in addition to the SAFINOUS score, the potential benefit of no-touch SVG harvesting should be considered.

Tomislav Kopjar, MD, PhD
Hrvoje Gasparovic, MD, PhD
Bojan Biocina, MD, PhD
Department of Cardiac Surgery
University of Zagreb School of Medicine and University
Hospital Center Zagreb
Zagreb, Croatia

References

1. Antonopoulos AS, Odutayo A, Oikonomou EK, Trivella M, Petrou M, Collins GS, et al. Development of a risk score for early saphenous vein graft failure: an individual patient data meta-analysis. *J Thorac Cardiovasc Surg.* 2020;160:102-12.
2. Kodja K, Patel S, Weber MP, Luc JGY, Choi JH, Maynes EJ, et al. Graft patency after open versus endoscopic saphenous vein harvest in coronary artery bypass grafting surgery: a systematic review and meta-analysis. *Ann Cardiothorac Surg.* 2018;7:586-97.
3. Kopjar T, Dashwood MR. Endoscopic versus "no-touch" saphenous vein harvesting for coronary artery bypass grafting. *Angiology.* 2016;67:121-32.
4. Krishnamoorthy B, Critchley WR, Thompson AJ, Payne K, Morris J, Venkateswaran RV, et al. Study comparing vein integrity and clinical outcomes in open vein harvesting and 2 types of endoscopic vein harvesting for coronary artery bypass grafting. *Circulation.* 2017;136:1688-702.
5. Samano N, Geijer H, Liden M, Fremes S, Bodin L, Souza D. The no-touch saphenous vein for coronary artery bypass grafting maintains a patency, after 16 years, comparable to the left internal thoracic artery: a randomized trial. *J Thorac Cardiovasc Surg.* 2015;150:880-8.
6. Samano N, Souza D, Botelho Pinheiro B, Kopjar T, Dashwood M. Twenty-five years of no-touch saphenous vein harvesting for coronary artery bypass grafting: structural observations and impact on graft performance. *Brazil J Cardiovasc Surg.* 2020;35:91-100.
7. Sousa-Uva M, Neumann FJ, Ahlsson A, Alfonso F, Banning AP, Benedetto U, et al. 2018 ESC/EACTS guidelines on myocardial revascularization. *Eur J Cardiothorac Surg.* 2019;55:4-90.

<https://doi.org/10.1016/j.jtcvs.2020.03.155>

REPLY: THE NO-TOUCH SAPHENOUS VEIN: INCREASED PATENCY, BUT AT WHAT RISK?

Reply to the Editor:

I read with interest the letter by Kopjar and colleagues¹ regarding the role of the no-touch saphenous vein in coronary bypass surgery. There is no doubt that minimization of endothelial damage is key to the long-term patency rate of bypass conduit, and

The authors reported no conflicts of interest.

The *Journal* policy requires editors and reviewers to disclose conflicts of interest and to decline handling or reviewing manuscripts for which they may have a conflict of interest. The editors and reviewers of this article have no conflicts of interest.

in fact small randomized trials have reported excellent angiographic outcomes for the no-touch saphenous vein.² On the other hand, the no-touch harvesting technique is associated with a higher risk of surgical site complications,² an issue particularly relevant for high-risk patients such as those with diabetes or obesity. Until we have clear data to understand the risk-to-benefit ratio between improved patency rate and increased risk of surgical complications with the no-touch saphenous vein, alternative conduits (the radial artery in particular) should be preferred.

Mario Gaudino, MD
Department of Cardiothoracic Surgery
Weill Cornell Medicine
New York City, NY

References

1. Kopjar T, Gasparovic H, Biocina B. The no-touch saphenous vein should be considered in a risk score of vein graft failure. *J Thorac Cardiovasc Surg.* 2020;160:e1-2.
2. Deb S, Singh SK, de Souza D, Chu MWA, Whitlock R, Meyer SR, et al. SUPERIOR SVG: no touch saphenous harvesting to improve patency following coronary bypass grafting (a multi-centre randomized control trial, NCT01047449). *J Cardiothorac Surg.* 2019;14:85.

<https://doi.org/10.1016/j.jtcvs.2020.03.154>

REPLY FROM AUTHORS: VEIN GRAFT BIOLOGY AND THE RISK OF GRAFT OCCLUSION

Reply to the Editor:

We thank Kopjar and colleagues¹ for their letter on the Saphenous Vein Graft Failure—An Outcomes Study in Coronary Artery Bypass Grafting (SAFINOUS-CABG) study,² and for giving us the opportunity to comment on the importance of vein harvesting for graft patency.

The SAFINOUS-CABG study² was an international collaboration between cardiothoracic centers with shared individual patient data from patients undergoing CABG and available angiographic data within the first year after surgery, and as such it was the first systematic effort to develop a comprehensive prediction model for early

FIGURE 1. Vein graft endothelial function and risk of occlusion after coronary artery bypass grafting (CABG). A, Occluded saphenous vein graft (SVG) post-CABG had significantly lower ex vivo vasorelaxations in response to acetylcholine (Ach) at the time of the operation. B, Predicted probability for SVG occlusion after adjustment for confounders based on the vasorelaxation response to Ach. OR, Odds ratio; CI, confidence interval; SD, standard deviation.

saphenous vein graft (SVG) occlusion. The developed SAFINOUS risk score² is composed of clinical, laboratory, and operative factors and predicts the risk for early SVG occlusion within the first year post-CABG.

In the SAFINOUS-CABG cohort, the use of endoscopic vein harvesting (EVH) was a protective factor against SVG occlusion.² The effects of EVH on graft patency are debatable; for example a recent meta-analysis showed that EVH may indeed be associated with higher rates of graft occlusion compared with open vein harvesting.³ We acknowledge that this may be related to the training and experience of operators, and therefore our findings on EVH may not be directly extrapolated to every clinical center. External validations of the SAFINOUS score from independent researchers would be welcome.

We totally agree with Kopjar and colleagues¹ on the value of a no-touch technique for SVG harvesting, which has been associated with reduced endothelial injury, and possibly also with a lower risk for early graft occlusion.⁴ In the Oxford cohort of the SAFINOUS-CABG study, SVG samples from all patients were used to assess vein redox state and endothelial function in organ bath experiments (as described in our previous studies).⁵ We observed that occluded SVG post-CABG had significantly lower ex vivo vasorelaxations in response to acetylcholine at the time of the operation (Figure 1, A). In logistic regression after adjustment for confounders, the vein endothelium-dependent vasodilation was independently associated with SVG occlusion (odds ratio, 0.177; 95% confidence interval, 0.058-0.545; $P = .003$ for percent acetylcholine-induced vasorelaxation above the median). The predicted probability for SVG occlusion after adjustment for confounders based on the vasorelaxation response to acetylcholine is

shown in Figure 1, B. This information on graft endothelial function was not uniformly available in the SAFINOUS-CABG cohort² and could not be included in the developed SAFINOUS score. However, these are important preliminary observations; they support the significance of vein biology for SVG patency and corroborate the arguments of Kopjar and colleagues¹ for the use of a no-touch harvesting technique to preserve vein integrity and graft endothelial function. Moreover, these observations highlight the role of vein endothelial function as a therapeutic target for the prevention of graft failure, which should be further explored in appropriately designed studies.

Future iterations of the SAFINOUS score could include such biological predictors for the risk of early graft occlusion. Nonetheless information on the surgical harvesting technique (eg, no-touch and graft probing) may also be good surrogates of vein integrity and can be more easily incorporated in a clinical score for the stratification of patients undergoing CABG for the risk of SVG failure.

Alexios S. Antonopoulos, MD, PhD^a

Attila Kardos, MD, PhD, FESC^b

Charalambos Antoniades, MD, PhD, FESC^a

^aDivision of Cardiovascular Medicine

Radcliffe Department of Medicine

University of Oxford

Oxford, United Kingdom

^bDivision of Cardiovascular Medicine

Milton Keynes University Hospital

NHS Foundation Trust

Milton Keynes, United Kingdom

The authors reported no conflicts of interest.

The *Journal* policy requires editors and reviewers to disclose conflicts of interest and to decline handling or reviewing manuscripts for which they may have a conflict of interest. The editors and reviewers of this article have no conflicts of interest.

References

1. Kopjar T, Gasparovic H, Biocina B. The no-touch saphenous vein should be considered in a risk score of vein graft failure. *J Thorac Cardiovasc Surg.* 2020; 160:e1-2.
2. Antonopoulos AS, Odutayo A, Oikonomou EK, Trivella M, Petrou M, Collins GS, et al. Development of a risk score for early saphenous vein graft failure: an individual patient data meta-analysis. *J Thorac Cardiovasc Surg.* 2020;160: 102-12.
3. Kodia K, Patel S, Weber MP, Luc JGY, Choi JH, Maynes EJ, et al. Graft patency after open versus endoscopic saphenous vein harvest in coronary artery bypass grafting surgery: a systematic review and meta-analysis. *Ann Cardiothorac Surg.* 2018;7:586-97.
4. Deb S, Singh SK, de Souza D, Chu MWA, Whitlock R, Meyer SR, et al. SUPERIOR SVG: no touch saphenous harvesting to improve patency following coronary bypass grafting (a multi-centre randomized control trial, NCT01047449). *J Cardiothorac Surg.* 2019;14:85.
5. Antoniadis C, Bakogiannis C, Tousoulis D, Reilly S, Zhang MH, Paschalis A, et al. Preoperative atorvastatin treatment in CABG patients rapidly improves vein graft redox state by inhibition of Rac1 and NADPH-oxidase activity. *Circulation.* 2010;122(11 Suppl):S66-73.

<https://doi.org/10.1016/j.jtcvs.2020.04.048>