the RV on LV performance. This compelling work sparks some new questions that deserve attention.

References

1. Gatzoulis MA, Balaji S, Webber SA, Siu SC, Hokanson JS, Poile C, et al. Risk factors for arrhythmia and sudden cardiac death late after repair of tetralogy of Fallot: a multicentre study. *Lancet*. 2000;356:975-81.
2. Geva T. Is MRI the preferred method for evaluating right ventricular size and function in patients with congenital heart disease? MRI is the preferred method for evaluating right ventricular size and function in patients with congenital heart disease. *Circ Cardiovasc Imaging*. 2014;7:190-7.
3. Valente AM, Gauvreau K, Assenza GE, Babu-Narayan SV, Schreier J, Gatzoulis MA, et al. Contemporary predictors of death and sustained ventricular tachycardia in patients with repaired tetralogy of Fallot enrolled in the INDICATOR cohort. *Heart*. 2014;100:247-53.
4. Brookes C, Ravn H, White P, Moeldrup U, Oldershaw P, Redington A. Acute right ventricular dilatation in response to ischemia significantly impairs left ventricular systolic performance. *Circulation*. 1999;100:761-7.
5. Stankovic Z, Allen BD, Garcia J, Jarvis KB, Markl M. 4D flow imaging with MRI. *Cardiovasc Diagn Ther*. 2014;4:173-92.
6. Schäfer M, Browne LP, Jagers J, Barker AJ, Morgan GJ, Ivy DD, et al. Abnormal left ventricular flow organization following repair of tetralogy of Fallot. *J Thorac Cardiovasc Surg*. 2020;160:1008-15.

See Article page 1008.

Commentary: Does right ventricular dysfunction cause left ventricular dysfunction in tetralogy of Fallot? The quest continues

William M. DeCampi, MD, PhD

Schafer and colleagues¹ present results of an analysis of left intraventricular flow dynamics (LVICF) in a group of 11 patients with repaired tetralogy of Fallot (TOF) with chronic pulmonary regurgitation (PR) and moderate increase in right ventricular end systolic volume index and right ventricular end diastolic volume index. They found that, compared with an age-matched control group of so-called normal subjects, the TOF group had abnormal flow patterns suggestive of left ventricle (LV) diastolic dysfunction. They showed that LVICF correlated with right ventricular end systolic volume index and end diastolic volume index and RV ejection fraction, suggesting the possibility of a causal relationship between abnormal RV function (in turn suggested to be due to PR) and abnormal LV diastolic function.

Tissue Doppler can detect LV dysfunction in TOF. MRI-based flow partitioning can detect this, too.

CENTRAL MESSAGE

Left ventricular dysfunction is common after tetralogy of Fallot repair. The challenge is to definitively determine the role of the right ventricle in causing the dysfunction.

Because these patients otherwise did not have conventional indications for pulmonary valve replacement, the authors suggested that LVICF measurements might be used to time earlier intervention on PR.

The authors used a method based on magnetic resonance imaging flow tracking developed by Eriksson and colleagues² to look at patterns of flow in the LV during diastole. These patterns are determined by the functional behavior of myocardial relaxation and recoil during diastole and in normal hearts optimize the efficiency of subsequent ejection. In effect, these patterns optimize alignment of fluid momentum to be efficiently additive to the momentum imparted to the fluid during systole. Any underlying cause of abnormal LV diastolic function will adversely

From the Department of Clinical Sciences, University of Central Florida College of Medicine, and Division of Cardiothoracic Surgery, Arnold Palmer Hospital for Children, Orlando, Fla.

Disclosures: The author reported no conflicts of interest.

The *Journal* policy requires editors and reviewers to disclose conflicts of interest and to decline handling or reviewing manuscripts for which they may have a conflict of interest. The editors and reviewers of this article have no conflicts of interest.

Received for publication Nov 20, 2019; revisions received Nov 20, 2019; accepted for publication Nov 21, 2019; available ahead of print July 3, 2020.

Address for reprints: William M. DeCampi, MD, PhD, The Heart Center, Arnold Palmer Hospital for Children, 80 W Miller St, 2nd Floor, Orlando, FL 32813 (E-mail: William.decampi@orlandohealth.com).

J Thorac Cardiovasc Surg 2020;160:1017-8

0022-5223/\$36.00

Copyright © 2020 by The American Association for Thoracic Surgery

<https://doi.org/10.1016/j.jtcvs.2019.11.153>

influence this flow pattern and subsequent systolic efficiency.

It has long been speculated that abnormal RV function may be a cause of abnormal LV diastolic function in repaired TOF. This causal relationship is suggested, but not proven by associations of indices of RV function with those of LV function. A definite mechanism has not been determined. For example, in 2012 Friedberg and colleagues³ showed that lower LV early diastolic radial and circumferential strain rates (by tissue Doppler echocardiography) were associated with greater RV volumes and greater PR in a study of 53 patients with repaired TOF and moderately increased RV volumes.³

In the present study, associations between LVICF and RV function indices are the basis for the suggestion that RV and LV dysfunction may be causally related. The calculated correlations should be carefully interpreted. The correlation analyses were performed using the combined cohort (n = 22). Although TOF patients will be scattered somewhat about the plot, the normal subjects will tend to cluster in 1 corner of the plot, almost surely yielding a significant linear correlation. Additional information could be obtained by computing the correlation within the TOF group alone (albeit it with larger sample size). Another problem (mentioned by the authors) is that the choice of normal subjects for the control group did not permit delineation of

factors that could influence LV and RV function independently. Examples include the nature of the congenital heart defect, effects of surgical intervention, and conduction abnormalities. Better control groups might consist of patients who underwent a successful valve-sparing TOF repair, or patients who underwent ventricular septal defect repair.

The authors should be applauded for continuing to aggressively pursue sensitive and reliable indicators of ventricular dysfunction after repair of congenital heart defects—TOF in particular. Our quest to circumvent LV dysfunction simply by changing something about the RV, however, must rely on well-controlled studies that establish causality and a mechanism the basis on which we can target intervention. Intuitively, we believe the target is a competent and durable pulmonary valve placed early in life. It might be more complicated than that.

References

1. Schafer M, Browne L, Jagers J, Barker A, Morgan G, Ivy D, et al. Abnormal left ventricular flow organization following repair of tetralogy of Fallot. *J Thorac Cardiovasc Surg.* 2020;160:1008-15.
2. Eriksson J, Dyverfeldt P, Engvall J, Bolger A, Ebbers T, Carhall C. Quantification of presystolic blood flow organization and energetics in the human left ventricle. *Am J Physiol Heart Circ Physiol.* 2011;300:H2135-41.
3. Friedberg M, Fernandes F, Roche L, Grosse-Wortman L, Manlhiot C, Fackoury C, et al. Impaired right and left ventricular diastolic myocardial mechanics and filling in asymptomatic children and adolescents after repair of tetralogy of Fallot. *Eur Heart J Cardiovasc Imaging.* 2012;13:905-13.