suit the new technology, we should examine if the thoughtful eye of an experienced aortic surgeon isn't already meeting the standard.

References

1. Elefteriades JA, Mukherjee SK, Mojibian H. Discrepancies in measurement of the thoracic aorta: JACC review topic of the week. *J Am Coll Cardiol*. 2020;76:201-17.
2. Goldstein SA, Evangelista A, Abbata S, Arai A, Asch FM, Badano LP, et al. Multimodality imaging of diseases of the thoracic aorta in adults: from the American Society of Echocardiography and the European Association of Cardiovascular Imaging: endorsed by the Society of Cardiovascular Computed Tomography and Society for Cardiovascular Magnetic Resonance. *J Am Soc Echocardiogr*. 2015;28:119-82.
3. Freeman LA, Young PM, Foley TA, Williamson EE, Bruce CJ, Greason KL. CT and MRI assessment of the aortic root and ascending aorta. *AJR Am J Roentgenol*. 2013;200:W581-92.
4. Vinholo TF, Zafar MA, Papanikolaou D, Chung J, Ellauzi H, Ziganshin BA, et al. Direct measurement of ascending aortic diameter by intraoperative caliper assessment. *J Thorac Cardiovasc Surg*. 2021;161:e143-6.
5. Ziganshin BA, Zafar MA, Elefteriades JA. Descending threshold for ascending aortic aneurysmectomy: is it time for a "left-shift" in guidelines? *J Thorac Cardiovasc Surg*. 2019;157:37-42.

See Article page e143.

Commentary: Measure twice, cut once

Jonathan C. Hong, MD, MHS, and
Joseph S. Coselli, MD

Jonathan C. Hong, MD, MHS, and Joseph S. Coselli, MD

The risk of aortic rupture and dissection is directly related to the diameter of the aorta according to Laplace's law. For more than 2 decades, the natural history studies from the Aortic Institute at Yale-New Haven Hospital have emphasized the increased incidence of aortic complications at larger aortic diameters.^{1,2} Diameter-based aortic thresholds, based on radiographic findings, are a critical consideration for surgical decision-making and form the basis of the clinical practice guidelines for surgical repair.³ For asymptomatic patients, the timing of operative repair is largely determined by preoperative imaging studies;

CENTRAL MESSAGE

Intraoperative direct caliper measurements of the ascending aorta correlate well with preoperative CT and TTE measurements at a high-volume aortic institution.

From the Division of Cardiothoracic Surgery, Michael E. DeBakey Department of Surgery, Baylor College of Medicine; Department of Cardiovascular Surgery, Texas Heart Institute; and CHI St Luke's-Baylor St. Luke's Medical Center, Houston, Tex.

Disclosures: Dr Coselli's work is partly supported by the Cullen Foundation. Dr Coselli consults for, receives royalties and a departmental educational grant from, and participates in clinical trials for Terumo Aortic; consults and participates in clinical trials for Medtronic, Inc, and W.L. Gore & Associates; and serves as a co-investigator for CytoSorbents. Dr Hong reported no conflicts of interest.

The *Journal* policy requires editors and reviewers to disclose conflicts of interest and to decline handling or reviewing manuscripts for which they may have a conflict of interest. The editors and reviewers of this article have no conflicts of interest.

Received for publication Aug 18, 2020; revisions received Aug 18, 2020; accepted for publication Aug 19, 2020; available ahead of print Aug 24, 2020.

Address for reprints: Joseph S. Coselli, MD, One Baylor Plaza, BCM 390, Houston, TX 77030 (E-mail: jcoselli@bcm.edu).

J Thorac Cardiovasc Surg 2021;161:e148-9

0022-5223/\$36.00

Copyright © 2020 by The American Association for Thoracic Surgery

<https://doi.org/10.1016/j.jtcvs.2020.08.059>

modalities include echocardiography (transthoracic [TTE] and transesophageal), computed tomography (CT), and magnetic resonance imaging. However, the ascending aorta can be challenging to image due to motion, oblique course, and variability of measurement techniques, as recently described by Elefteriades and coauthors.⁴

In this current issue of the *Journal*, Vinholo and coauthors⁵ from the Aortic Institute at Yale-New Haven Hospital compared intraoperative, direct caliper measurements to preoperative imaging studies, primarily CT and TTE measurements. In this study of 35 patients, preoperative measurements corresponded well with intraoperative measurements of the ascending aorta. Nuances of measurement included the exclusion of the aortic

wall by CT angiography, the inclusion of the aortic wall via caliper, and partial exclusion of the aortic wall by leading-edge-to-leading-edge techniques of TTE. Despite inherent differences between preoperative and intraoperative measurements, such as how the aortic wall was captured, the authors found preoperative radiographic measurements to be well correlated with intraoperative measurement. Notably, because the natural history of aortic disease is tied to preoperative imaging studies, it is reassuring that these radiographic techniques approximate the true aortic diameter as obtained through direct intraoperative measurement. However, the use of intraoperative measurements has limited value in scheduling patients for repair. In dynamic scenarios in which the patient is on the table for other reasons (ie, valvular concerns or to address coronary artery disease) and aortic disease presents unexpectedly, the surgeon may gain confidence knowing that intraoperative, direct measurement by caliper is highly correlated to established diameter-based thresholds of repair.

Vinholo and coauthors provide a key contribution in empirically examining the assumption that preoperative imaging correlates well to the true dimensions of the ascending aorta. Even so, the aortic dimensions were measured at a high-volume aortic surgery institution with well-defined protocols in preoperative imaging and may not be generalizable to all institutions. Elefteriades and coauthors⁴ recently described the various sources of discrepancy that occur within and between imaging modalities. Elsewhere, the transcatheter community has recently put forth an expert consensus on CT imaging in the context of transcatheter aortic valve replacement.⁶ Similarly, aortic surgeons should work together with

imaging specialists in standardizing how we measure the aorta, thereby strengthening the foundation upon which our surgical decision-making rests. In addition, literature describing aortic repair should aim to be more precise when we describe diameters and include the measurement technique. For the foreseeable future until we have better ways to assess the strength, thickness, and integrity of the aortic wall, measurements of the aortic diameter will continue to serve as the hinge point for repair.

References

1. Coady MA, Rizzo JA, Hammond GL, Mandapati D, Darr U, Kopf GS, et al. What is the appropriate size criterion for resection of thoracic aortic aneurysms? *J Thorac Cardiovasc Surg*. 1997;113:476-91; discussion 89-91.
2. Chau KH, Elefteriades JA. Natural history of thoracic aortic aneurysms: size matters, plus moving beyond size. *Prog Cardiovasc Dis*. 2013;56:74-80.
3. Hiratzka LF, Bakris GL, Beckman JA, et al. 2010 ACCF/AHA/AATS/ACR/ASA/SCA/SCAI/SIR/STS/SVM guidelines for the diagnosis and management of patients with thoracic aortic disease. A report of the American College of Cardiology Foundation/American Heart Association task force on practice guidelines, American Association for Thoracic Surgery, American College of Radiology, American Stroke Association, Society of Cardiovascular Anesthesiologists, Society for Cardiovascular Angiography and Interventions, Society of Interventional Radiology, Society of Thoracic Surgeons, and Society for Vascular Medicine. *J Am Coll Cardiol*. 2010;55:e27-129.
4. Elefteriades JA, Mukherjee SK, Mojibian H. Discrepancies in measurement of the thoracic aorta: JACC review topic of the week. *J Am Coll Cardiol*. 2020;76:201-17.
5. Vinholo TF, Zafar MA, Papanikolaou D, Chung J, Ellauzi H, Ziganshin BA, et al. Direct measurement of ascending aortic diameter by intraoperative caliper assessment. *J Thorac Cardiovasc Surg*. 2021;161:e143-6.
6. Blanke P, Weir-McCall JR, Achenbach S, Delgado V, Hausleiter J, Jilaihawi H, et al. Computed tomography imaging in the context of transcatheter aortic valve implantation (TAVI)/transcatheter aortic valve replacement (TAVR): an expert consensus document of the Society of Cardiovascular Computed Tomography. *JACC Cardiovasc Imaging*. 2019;12:1-24.