and refinement as needed, may help surgical teams get a better grasp of functional status for patients being considered for esophagectomy.

References

1. Tang A, Raja S, Rappaport J, Raymond D, Sudarshan M, Bribrisco A, et al. Looking beyond the eyeball test: a novel vitality index to predict recovery after esophagectomy. *J Thorac Cardiovasc Surg*. 2021;161:822-32.e6.
2. Ferguson MK, Thompson K, Huisin-Scheetz M, Farnan J, Hemmerich JA, Slawinski K, et al. Thoracic surgeons' perception of frail behavior in videos of standardized patients. *PLoS One*. 2014;9:e98654.
3. Holeman TA, Peacock J, Beckstrom JL, Brooke BS. Patient-surgeon agreement in assessment of frailty, physical function, and social activity. *J Surg Res*. 2020; 256:368-73.
4. Mrdutt MM, Papaconstantinou HT, Robinson BD, Bird ET, Isbell CL. Preoperative frailty and surgical outcomes across diverse surgical subspecialties in a large health care system. *J Am Coll Surg*. 2019;228:482-90.
5. Hodari A, Hammoud ZT, Borgi JF, Tsiouris A, Rubinfeld IS. Assessment of morbidity and mortality after esophagectomy using a modified frailty index. *Ann Thorac Surg*. 2013;96:1240-5.
6. Shah R, Attwood K, Arya S, Hall DE, Johanning JM, Gabriel E, et al. Association of frailty with failure to rescue after low-risk and high-risk inpatient surgery. *JAMA Surg*. 2018;153:e180214.
7. Hirpara DH, Kidane B, Rogalla P, Cypel M, de Perrot M, Keshavjee S, et al. Frailty assessment prior to thoracic surgery for lung or esophageal cancer: a feasibility study. *Support Care Cancer*. 2019;27:1535-40.
8. Sandini M, Pinotti E, Persico I, Picone D, Bellelli G, Gianotti L. Systematic review and meta-analysis of frailty as a predictor of morbidity and mortality after major abdominal surgery. *BJS Open*. 2017;1:128-37.
9. Sheetz KH, Zhao L, Holcombe SA, Wang SC, Reddy RM, Lin J, et al. Decreased core muscle size is associated with worse patient survival following esophagectomy for cancer. *Dis Esoph*. 2013;26:716-22.
10. Guinan EM, Doyle SL, Bennett AE, O'Neill L, Gannon J, Elliott JA, et al. Sarcopenia during neoadjuvant therapy for oesophageal cancer: characterising the impact on muscle strength and physical performance. *Support Care Cancer*. 2018;26:1569-76.

See Article page 822.

Commentary: Surgical risk assessment in 2020: Is a handshake and a walking test really the best we've got?

Ernest G. Chan, MD, MPH,
Chigozirim N. Ekeke, MD, and
James D. Luketich, MD

Minimally invasive approaches to many disease processes, including esophageal cancer, can help lower the risk of morbidity and mortality.¹ Accurate preoperative assessment is a key component in stratifying patients who present to clinic for possible esophagectomy. These tools allow both medical oncologists and general thoracic surgeons the ability to predict the outcomes of various treatment plans,

Ernest G. Chan, MD, MPH, Chigozirim N. Ekeke, MD, and James D. Luketich, MD

CENTRAL MESSAGE

There is a need for better pre-operative assessment in patients undergoing esophagectomy. The Esophagectomy Vitality Index is a novel system that assesses physical status and fitness in these patients.

From the Department of Cardiothoracic Surgery, University of Pittsburgh School of Medicine and the University of Pittsburgh Medical Center, Pittsburgh, Pa.

Disclosures: Dr Luketich owns stock in Intuitive Surgical and Express Scripts. All other authors reported no conflicts of interest.

The *Journal* policy requires editors and reviewers to disclose conflicts of interest and to decline handling or reviewing manuscripts for which they may have a conflict of interest. The editors and reviewers of this article have no conflicts of interest.

Received for publication Nov 11, 2020; revisions received Nov 11, 2020; accepted for publication Nov 12, 2020; available ahead of print Nov 20, 2020.

Address for reprints: James D. Luketich, MD, FACS, Department of Cardiothoracic Surgery, University of Pittsburgh, School of Medicine, C800 PUH, 200 Lothrop St, Pittsburgh, PA 15213 (E-mail: luketichjd@upmc.edu).

J Thorac Cardiovasc Surg 2021;161:834-5

0022-5223/\$36.00

Copyright © 2020 Published by Elsevier Inc. on behalf of The American Association for Thoracic Surgery

<http://dx.doi.org/10.1016/j.jtcvs.2020.11.040>

including multimodal therapy and esophagectomy, and identify nonsurgical candidates. An accurate predictive assessment might also help us decide when a period of physical rehabilitation and medical “tune-up” might be considered when the initial risk assessment appears concerning. While the literature has presented many methods for assessing surgical candidacy,²⁻⁶ there continues to be a need to improve our preoperative assessment tools, since esophagectomy continues to be associated with major morbidity and mortality even in the era of minimally invasive esophagectomy.⁷

Tang and colleagues⁸ present their novel Esophagectomy Vitality Index to assess the surgical candidacy of patients with esophageal cancer. Several characteristics of this novel index help differentiate the Esophagectomy Vitality Index from other physiological indices. The simplistic-yet-powerful design of the index in question should be highlighted. This novel vitality index consists of 4 items that test a patient's performance parameters and preoperative functional status. These 4 items are performed during the clinic visit (grip strength, 30-second chair sit-stands, 6-minute walk) or obtained through computed tomography imaging (psoas muscle area), lending to the accessibility of performing this index.⁸

The Esophagectomy Vitality Index involves identifying a patient's physical status and fitness just before surgery and therefore may offer the best real-time data. However, while the authors provide standardization in each component, there may be too much variability associated with each test. For example, lower-extremity length may introduce variability in difficulty with the 30-second chair sit-stand test. Similarly, hand size could cause differences in grip strength measures even on a calibrated dynamometer. In addition, there may be anatomic variations in muscle insertion and origin independent of height. Other important details could be further assessed in normal volunteers, such as the reproducibility of these functional variables. In addition, the variability of multiple attempts at a 30-second sit-stand by different observers was not entirely clear, and the potential for a period of physical therapy and rehabilitation with a follow-up index should the initial index score be concerning. All of these are opportunities to strengthen this index and prove that it is reproducible.

Despite these issues, the addition of the Esophagectomy Vitality Index will help us move away from the "eyeball"

test. The existing knowledge of quantifying "surgical fitness" will further improve the predictive capability of this index score. Indeed, the work presented by Tang and colleagues serves as a platform for improving how surgeons define and quantify "surgical fitness" as an objective measurement for surgical candidacy. We applaud Tang and his colleagues for this contribution to the literature and look forward to witnessing future validation and implementation in prospective, possibly randomized trials, and how this might impact on future esophagectomy practices.

References

1. Luketich JD, Pennathur A, Awais O, Levy RM, Keeley S, Shende M, et al. Outcomes after minimally invasive esophagectomy: review of over 1000 patients. *Ann Surg*. 2012;256:95-103.
2. Hu Y, McMurry TL, Stukenborg GJ, Kozower BD. Readmission predicts 90-day mortality after esophagectomy: analysis of surveillance, epidemiology, and end results registry linked to Medicare outcomes. *J Thorac Cardiovasc Surg*. 2015;150:1254-60.
3. Janowak CF, Blasberg JD, Taylor L, Maloney JD, Macke RA. The surgical Apgar score in esophagectomy. *J Thorac Cardiovasc Surg*. 2015;150:806-12.
4. Reeh M, Metze J, Uzunoglu FG, Nentwich M, Ghadban T, Wellner U, et al. The PER (Preoperative Esophagectomy Risk) score: a simple risk score to predict short-term and long-term outcome in patients with surgically treated esophageal cancer. *Medicine (Baltimore)*. 2016;95:e2724.
5. Van Veer H, Moons J, Darling G, Lerut T, Coosemans W, Waddell T, et al. Validation of a new approach for mortality risk assessment in esophagectomy for cancer based on age- and gender-corrected body mass index. *Eur J Cardiothorac Surg*. 2015;48:600-7.
6. Hodari A, Hammoud ZT, Borgi JF, Tsiouris A, Rubinfeld IS. Assessment of morbidity and mortality after esophagectomy using a modified frailty index. *Ann Thorac Surg*. 2013;96:1240-5.
7. Seder CW, Raymond D, Wright CD, Gaisert HA, Chang AC, Becker S, et al. The Society of Thoracic Surgeons general thoracic surgery database 2018 update on outcomes and quality. *Ann Thorac Surg*. 2018;105:1304-7.
8. Tang A, Ahmad U, Raja S, Rappaport J, Raymond DP, Sudarshan M, et al. Looking beyond the eyeball test: a novel vitality index to predict recovery after esophagectomy. *J Thorac Cardiovasc Surg*. 2021;161:822-32.e6.