See Article page 1155.

Commentary: Complication monitoring comes of age

Camille L. Hancock Friesen, MD

In their article in this issue of the *Journal*, Dorobantu and colleagues¹ report a large contemporary series evaluating the incidence of post–pediatric cardiac surgery unplanned reinterventions (uREs), the risk factors for uRE occurrence, and the impact of uRE occurrence on mortality.¹ Their findings will come as no surprise to anyone who has been following the emergence of complication monitoring and reporting in pediatric cardiac surgery. Five centers (representing approximately 50% of pediatric cardiac surgery cases performed in the United Kingdom) collaborated to capture uREs, defined as the occurrence of any of the following: cardiac reoperation, interventional catheterization, pacemaker implantation, or diaphragm plication. The authors prospectively collected complication data from 3090 index procedures (defined in this study as “30d surgical episodes”) performed in 2861 patients and reported on 153 uREs occurring in 146 patients (uRE rate, 4.7%). The incidence of the individual uREs was low and laudable; 2.9% for cardiac reoperation, 0.8% for interventional catheterization, 0.8% for permanent pacemaker, and 0.4% for diaphragm plication. Despite different definitions of “index procedure” and uRE, the rates in this report are comparable to reports from the Society of Thoracic Surgeons Congenital Heart Surgery Database.^{2–4} Higher-risk operations are associated with more complications, and more complications are associated with higher mortality, as has been reported previously.^{5,6} In this report, the authors used a stringent matching algorithm to generate 74 pairs of patients with uREs (cases) and patients with no uREs (controls) and report the novel finding of an ongoing subacute (at 6 months postoperation) hazard for mortality

Camille L. Hancock Friesen, MD

CENTRAL MESSAGE

Unplanned reinterventions (uREs) carry an independent ongoing subacute hazard for mortality in pediatric cardiac surgery patients. uREs can be used as part of a quality metric and as a target for improving outcomes.

attributable in patients with uRE (12.2% in cases vs 1.4% in controls; $P = .02$), independent of other risk factors.

Although the authors make it look easy, the analysis and presentation of complication data are anything but. Concise definitions of complications are mandatory. The degree of surveillance (ie, how hard one looks for a complication) impacts the incidence, and here the prospective nature of the study is a clear strength of the report. In an exemplary model of coordination and collaboration, the multidisciplinary multisite team assembled and articulated a priori definitions and ultimately developed a system for prospectively capturing complications.

One inherent assumption of this type of analysis is that complication occurrence is linear; that is, complications occur independent of one another. Dorobantu and colleagues do a heroic job attempting to identify which patient and procedure factors affect survival after uREs and use novel mapping graphics to illustrate the complexity of complication clustering and frequency of occurrence. As big data, artificial intelligence, and machine learning become part of our ecosystem, we can expect further explanation of the complexities of complication occurrence.⁷

As a discipline, we can now name and count complications, and indeed we have had explicit definitions of complications for more than a decade.⁸ We know that regardless of the volume of a program, nearly one-half of heart surgery

From the Division of Pediatric Cardiothoracic Surgery, Children's Health Dallas and University of Texas Southwestern, Dallas, Tex.

Disclosures: The author reported no conflicts of interest.

The *Journal* policy requires editors and reviewers to disclose conflicts of interest and to decline handling or reviewing manuscripts for which they may have a conflict of interest. The editors and reviewers of this article have no conflicts of interest.

Received for publication Nov 13, 2020; revisions received Nov 13, 2020; accepted for publication Nov 16, 2020; available ahead of print Nov 23, 2020.

Address for reprints: Camille L. Hancock Friesen, MD, Children's Health Dallas, 1935 Medical District Drive, Dallas, TX 75235 (E-mail: Camille.HancockFriesen@UTSouthwestern.edu).

J Thorac Cardiovasc Surg 2021;161:1167–8

0022-5223/\$0.00

Published by Elsevier Inc. on behalf of The American Association for Thoracic Surgery

<https://doi.org/10.1016/j.jtcvs.2020.11.052>

patients will experience at least 1 complication.^{5,9} We also know that a key contributor to reduced mortality is timely recognition and management of complications, sometimes termed “rescuing.”⁹ With this and other recent reports, we are embracing complication analysis as part of the assessment of quality of care.^{10,11} Moving upstream of complications to prevent their occurrence will be the next worthy challenge in this dynamic and evolving field. The description of the magnitude and consequences of any problem is the essential foundation to the fundamental goal of reducing (or eliminating) that problem, and Dorobantu and colleagues have provided such a foundation with this work.

References

1. Dorobantu DM, Ridout D, Brown KL, Rodrigues W, Sharabiani MTA, Pagel C, et al. Factors associated with unplanned reinterventions and their relation to early mortality after pediatric cardiac surgery. *J Thorac Cardiovasc Surg.* 2021;161:1155-66.e9.
2. Costello JM, Mongé MC, Hill KD, Kim S, Pasquali SK, Yerokun BA, et al. Associations between unplanned cardiac reinterventions and outcomes after pediatric cardiac operations. *Ann Thorac Surg.* 2018;105:1255-63.
3. Thibault D, Wallace AS, Jacobs ML, Hornik CP, Costello JM, Fleming GF, et al. Postoperative transcatheter interventions in children undergoing congenital heart surgery. *Circ Cardiovasc Interv.* 2019;12:e007853.
4. O'Brien SM, Jacobs JP, Shahian DM, Jacobs ML, Gaynor JW, Romano JC, et al. Development of a congenital heart surgery composite quality metric: part 2-analytic methods. *Ann Thorac Surg.* 2019;107:590-6.
5. Belliveau D, Burton HJ, O'Blens SB, Warren AE, Hancock Friesen CL. Real-time complication monitoring in pediatric cardiac surgery. *Ann Thorac Surg.* 2012;94:1596-602.
6. Brown KL, Ridout D, Pagel C, Wray J, Anderson D, Barron DJ, et al. Incidence and risk factors for important early morbidities associated with pediatric cardiac surgery in a UK population. *J Thorac Cardiovasc Surg.* 2019;158:1185-96.e7.
7. Bihorac A, Ozrazgat-Baslanti T, Ebadi A, Motaei EA, Madkour M, Pardalos PM, et al. MySurgeryRisk: development and validation of a machine-learning risk algorithm for major complications and death after surgery. *Ann Surg.* 2019;269:652-62.
8. Jacobs JP. Introduction—databases and the assessment of complications associated with the treatment of patients with congenital cardiac disease. *Cardiol Young.* 2008;18(Suppl 2):1-37.
9. Pasquali SK, He X, Jacobs JP, Jacobs ML, O'Brien SM, Gaynor JW. Evaluation of failure to rescue as a quality metric in pediatric heart surgery: an analysis of the STS Congenital Heart Surgery Database. *Ann Thorac Surg.* 2012;94:573-80.
10. Pasquali S, Shahian DM, O'Brien SM, Jacobs ML, Gaynor JW, Romano JC, et al. Development of a congenital heart surgery composite quality metric, part 1: conceptual framework. *Ann Thorac Surg.* 2019;107:583-9.
11. Pasquali SK, Thibault D, O'Brien SM, Jacobs JP, Gaynor JW, Romano JC, et al. National variation in congenital heart surgery outcomes. *Circulation.* 2020;142:1351-60.

See Article page 1155.

Commentary: Unplanned reinterventions in pediatric cardiac surgery: Second time's a charm?

T. Konrad Rajab, MD, and
Minoo N. Kavarana, MD, FACS

Unplanned reinterventions are important for a number of reasons. First and foremost, they may result from deficiencies in the quality of clinical care that may be improved.

From the Section of Pediatric Cardiothoracic Surgery, Medical University of South Carolina, Charleston, SC.

Disclosures: The authors reported no conflicts of interest.

The *Journal* policy requires editors and reviewers to disclose conflicts of interest and to decline handling or reviewing manuscripts for which they may have a conflict of interest. The editors and reviewers of this article have no conflicts of interest.

Received for publication Nov 13, 2020; revisions received Nov 13, 2020; accepted for publication Nov 16, 2020; available ahead of print Nov 24, 2020.

Address for reprints: Minoo N. Kavarana, MD, FACS, Section of Pediatric Cardiothoracic Surgery, 10 McClellan Banks Dr, Charleston, SC 29425 (E-mail: kavarana@musc.edu).

J Thorac Cardiovasc Surg 2021;161:1168-9

0022-5223/\$36.00

Copyright © 2020 by The American Association for Thoracic Surgery

<https://doi.org/10.1016/j.jtcvs.2020.11.061>

Unplanned reinterventions are associated with early mortality yet are a necessary evil.

CENTRAL MESSAGE

Unplanned reinterventions are associated with significantly increased early mortality. However, on occasion, they are a necessary evil to rescue patients from complications.

Second, unplanned reinterventions are adverse outcomes for patients since each procedure carries additional risks for morbidity and mortality. Further, unplanned