See Article page 1287.

Commentary: Vita ex machina—life from the machine

Hellmuth R. Muller Moran, MD,
Michael H. Yamashita, MDCM, MPH, FRCSC, and
Rakesh C. Arora, MD, PhD, FRCSC

It is no secret that the use of extracorporeal life support (ECLS) has increased dramatically. According to the Extracorporeal Life Support Organization registry, 463 centers performed 15,875 ECLS runs in 2019 compared with 115 centers that performed 1743 ECLS runs in 2000.¹ Several documents have been produced to guide the care of patients on ECLS on diverse topics from percutaneous mechanical circulatory support² to extracorporeal cardiopulmonary resuscitation,³ blood transfusion,⁴ and most recently the Coronavirus Disease 2019 pandemic.⁵ These efforts represent critical junctures in the care of their respective populations, because without suitable guidance there can be no progress. Until now, however, there has been minimal guidance on the care of patients in postcardiotomy cardiogenic shock, which occurs commonly and in whom outcomes are generally poor without mechanical support.

The authors, representing the Extracorporeal Life Support Organization, European Association for Cardio-Thoracic Surgery, Society for Thoracic Surgery, and American Association for Thoracic Surgery, are to be congratulated for producing a comprehensive expert consensus document on postcardiotomy ECLS in adults.⁶ This document discusses many aspects of preoperative, during, and post-ECLS care. Reasonable, well thought-out recommendations are provided for a variety of commonly encountered and a number of less commonly encountered—but no less important—scenarios. Although the

Hellmuth R. Muller Moran, MD, Michael H. Yamashita, MDCM, MPH, FRCSC, and Rakesh C. Arora, MD, PhD, FRCSC

CENTRAL MESSAGE

Although stronger evidence and guidance are still needed in many areas, providers at all levels will benefit from this comprehensive handbook on postcardiotomy ECLS care.

procedure of initiating ECLS is often technically straightforward, it is in the multitude of perioperative possibilities, including reaching futility that the “art” of ECLS is truly found. Indeed, apart from the initial status at presentation, these are the elements that perhaps have the greatest bearing on a patient’s ultimate course. Individuals who are involved with ECLS at any level will benefit greatly by considering the many recommendations contained herein and applying them to their practice when able.

The difficulty with this document and with many areas of acute and critical cardiac surgical care is the relatively low level of evidence supporting most of the included statements. Although there is often a significant amount of anecdotal evidence on how best to manage these challenging patients, Level A evidence is sparse. This highlights the need for rigorous, prospective studies or randomized controlled trials in the population on ECLS. Acknowledging this can be difficult given these patients are critically ill and not easily recruited into randomized studies. Nonetheless, it remains incumbent on our field to perform well-designed studies in patients on ECLS and to report our findings. Only by these efforts will we be able to further improve the care that we provide standardized practices across the field and identify the areas most deserving of future initiatives.

ECLS in postcardiotomy patients has changed the field of cardiac surgery, yet there remain many unanswered questions regarding its use. This important *opus* serves as an essential document for our specialty, particularly with provision of the most up-to-date guidance for the use of valuable ECLS resources in the postcardiotomy patient. This

From the Division of Cardiac Surgery, Department of Surgery, Max Rady College of Medicine, University of Manitoba, Winnipeg, Manitoba, Canada; and Cardiac Sciences Program, St Boniface Hospital, Winnipeg, Manitoba, Canada.

Disclosures: R.C.A. has unrestricted grant from Pfizer Canada Inc, and has received honoraria from Abbott Nutrition and Edwards Lifesciences for work unrelated to this article. All other authors reported no conflicts of interest.

The *Journal* policy requires editors and reviewers to disclose conflicts of interest and to decline handling or reviewing manuscripts for which they may have a conflict of interest. The editors and reviewers of this article have no conflicts of interest.

Received for publication Nov 15, 2020; revisions received Nov 15, 2020; accepted for publication Nov 17, 2020; available ahead of print Nov 24, 2020.

Address for reprints: Rakesh C. Arora, MD, PhD, FRCSC, I.H. Asper Clinical Research Institute, CR3015 - 369 Tache Ave, Winnipeg, Manitoba R2H 2A6, Canada (E-mail: rakeshcarora@gmail.com).

J Thorac Cardiovasc Surg 2021;161:1333-4

0022-5223/\$36.00

Copyright © 2020 by The American Association for Thoracic Surgery

<https://doi.org/10.1016/j.jtcvs.2020.11.066>

document will doubtless be considered required reading and, along with the *vade mecum*—or handbook—included in the supplementary materials, an invaluable resource for providers of postcardiotomy ECLS care across the globe.

References

1. Extracorporeal Life Support Organization. ECLS Registry Report International Summary - July 2020. Ann Arbor, MI: Extracorporeal Life Support Organization; 2020.
2. Rihal CS, Naidu SS, Givertz MM, Szeto WY, Burke JA, Kapur NK, et al. 2015 SCAI/ACC/HFSA/STS clinical expert consensus statement on the use of percutaneous mechanical circulatory support devices in cardiovascular care: endorsed by the American Heart Association, the Cardiological Society of India, and Sociedad Latino Americ. *J Am Coll Cardiol*. 2015;65:e7-26.
3. Panchal AR, Berg KM, Hirsch KG, Kudenchuk PJ, Del Rios M, Cabañas JG, et al. 2019 American Heart Association focused update on advanced cardiovascular life support: use of advanced airways, vasopressors, and extracorporeal cardiopulmonary resuscitation during cardiac arrest: an update to the American Heart Association guidelines. *Circulation*. 2019;140:e881-94.
4. Singh G, Nahirmiak S, Arora R, Légaré J-F, Kanji HD, Dave Nagpal D, et al. Transfusion thresholds for adult respiratory extracorporeal life support: an expert consensus document. *Can J Cardiol*. 2020;36:1550-3.
5. Shekar K, Badulak J, Peek G, Boeken U, Dalton HJ, Lovkesh Arora L, et al. Extracorporeal life support organization coronavirus disease 2019 interim guidelines: a consensus document from an international group of interdisciplinary extracorporeal membrane oxygenation providers. *ASAIO J*. 2020;66:707-21.
6. Lorusso R, Whitman G, Milojevic M, Raffa G, McMullan DM, Boeken U, et al. 2020 EACTS/ELSO/STS/AATS expert consensus on post-cardiotomy extracorporeal life support in adult patients. *J Thorac Cardiovasc Surg*. 2021;161:1287-331.