an alternative but a sine qua non surgical time for completing the resection.

The anterior approach of Masaoka, Ito, and Yasumitsu¹³ affords a superb anterior operative view for resecting bronchogenic tumors involving the anterior thoracic inlet, and we used their approach once.¹⁴ However, we believe that the median sternotomy and anterior intercostal thoracotomy are not necessary for resecting apical tumors, because they do not add much more exposure but do increase postoperative complications (e.g., flail chest¹³) and morbidity, especially when the phrenic nerve needs to be resected. The median sternotomy yields a transverse exposure of the mediastinal and pulmonary structures. In our opinion, this gives a more theoretical than practical surgical view to apical lesions in that it cannot afford adequate exposure and manipulation of those apical lesions involving, for instance, the anterior arches of the first ribs and surrounding structures. By contrast, our cervical approach yields a direct open window to the entire aperture of the thoracic inlet once the internal half of the clavicle has been removed; the key success of our approach is this last step of the operation, which permits a direct view and surgical management of all tumors involving the thoracic inlet. Moreover, by making the horizontal incision either immediately below the clavicle or in the bed of the second or third intercostal muscles and resecting the related chest wall, we obtain the equivalent of an anterior thoracotomy. In this sense, 10 anterior, 2 middle, and 4 posterior apical tumors were radically resected through the cervical approach alone.

Drs. Niwa and Masaoka also recommend operation for bronchogenic tumors involving the sternum or ipsilateral supraclavicular lymph nodes. We might well speculate that in Western countries the indications for resecting bronchogenic tumors are somewhat more cautious and limited, particularly in terms of N status. At any rate, the technique described by Niwa and associates¹⁵ might be an option, but we believe it simply represents an anterior prolongation above the nipple of the classic posterior approach described by Shaw, Paulson, and Kee⁷ in 1961. This technique frees the entire scapular region but has been abandoned at our institution because of its limited exposure of the upper compartments of the thoracic inlet and its excessive morbidity.

Philippe Darteville, MD
Paolo Macchiarini, MD

Department of Thoracic and Vascular Surgery
and Heart-Lung Transplantation
Hôpital Marie-Lannelongue (Paris-Sud University)
133, Avenue de la Résistance
92350 Le Plessis Robinson, France

REFERENCES

1. Darteville P, Chapelier A, Macchiarini P, et al. Anterior transcervical-thoracic approach for radical resection of lung tumors invading the thoracic inlet. *J THORAC CARDIOVASC SURG* 1993;105:1025-34.
2. Masaoka A, Mizuno T, Shibata K, Yamakawa Y, Niwa H. Characterization of the apical invading lung cancers at different localization: emphasis of anterior apical tumor. *Nagoya Med J* 1987;32:137-46.
3. Pancoast HK. Superior sulcus tumors. *JAMA* 1932;99:1391-6.
4. Paulson DL. The importance of defining location and staging of superior pulmonary sulcus tumors. *Ann Thorac Surg* 1973;15:549-51.
5. Teixeira JP. Concerning the Pancoast tumor: What is the superior pulmonary sulcus? *Ann Thorac Surg* 1983;35:577-8.
6. Tobias JW. Síndrome apico-costovertebral-doloroso por tumor apical: su valor diagnóstico en el cáncer primitivo del pulmón. Thesis, Buenos Aires: Impresora mercatali, 1931.
7. Shaw RR, Paulson DL, Kee JL Jr. Treatment of the superior sulcus tumor by irradiation followed by resection. *Ann Surg* 1961;154:29-40.
8. Shahian DM, Wildford BN, Ellis FH Jr. Pancoast tumors: improved survival with preoperative and postoperative radiotherapy. *Ann Thorac Surg* 1987;43:32-8.
9. Paulson DL. Carcinomas in the superior pulmonary sulcus. *J THORAC CARDIOVASC SURG* 1975;70:1095-104.
10. Anderson TM, Moy PM, Holmes EC. Factors affecting survival in superior sulcus tumors. *J Clin Oncol* 1985;4:598-603.
11. Pancoast HK. Importance of careful roentgen-ray investigations of apical chest tumors. *JAMA* 1924;83:1407-11.
12. Darteville P, Macchiarini P. Technique of anterior transcervical-thoracic resection of apical lung tumors. In: Pearson FG, Deslauriers J, Ginsberg RJ, Hiebert CA, McKneally MFC, Urschel HC, eds. *Thoracic surgery*. New York: Churchill Livingstone, [In press].
13. Masaoka A, Ito Y, Yasumitsu T. Anterior approach for tumors of the superior sulcus. *J THORAC CARDIOVASC SURG* 1979;78:413-5.
14. Macchiarini P, Chapelier A, Monnet I, et al. Extended operations after induction therapy for T4 non-small cell lung cancer. *Ann Thorac Surg* [In press].
15. Niwa H, Masaoka A, Yamakawa Y, Fukui I, Kiriya M. Surgical therapy for apical invasive lung cancer: different approaches according to tumor location. *Lung Cancer* 1993;10:63-71.

Late management of esophageal perforation

To The Editor:

I read with interest "Management of Delayed Esophageal Perforation" by Salo and others in the December 1993 issue of the JOURNAL.¹ How should this be managed, asked the authors: "Esophagectomy or primary repair?"

To answer this question, they report their experience. Among 19 patients treated with primary repair, 13 died, which constitutes a mortality of 68.4%. The mortality for the patients in the esophagectomy group was lower, 13%, but accompanied by high morbidity because of the magnitude of the operation and long-term care necessary for esophagectomy and later reconstitution of the gastrointestinal tract. The authors recognize that the lethal factor in this group of patients is mediastinal sepsis: "Insufficient control of the mediastinal sepsis is the principal cause of death in patients with delayed esophageal perforation."

A third alternative not mentioned by the authors of this paper directs therapy directly to the treatment of mediastinitis itself. In my experience and that of others, irrigation of the mediasti-

num through the esophageal opening has been rewarded with complete resolution of mediastinitis.² An extra benefit has been the spontaneous healing and closure of the esophageal opening.

If the authors had used transesophageal irrigation on their patients with a failed esophageal closure, it is possible that better results would have been obtained. I continue to recommend transesophageal irrigation in patients with mediastinitis caused by esophageal perforation as the best option for this serious condition.

*Gil Hauer Santos, MD
Thoracic Surgery
Albert Einstein College of Medicine
Bronx, NY 10461*

REFERENCES

1. Salo JA, Isolauri JO, Heikkilä LJ, et al. Management of delayed esophageal perforation with mediastinal sepsis. *J THORAC CARDIOVASC SURG* 1993;106:1088-91.
2. Santos GH, Frater RW. Transesophageal irrigation for the treatment of mediastinitis produced by esophageal rupture. *J THORAC CARDIOVASC SURG* 1986;91:57-62.

Reply to the Editor:

Drs. Santos and Frater¹ reported a very interesting technique to treat patients with mediastinitis caused by esophageal perforation. They irrigated the mediastinum through the perforation in the esophagus. Profuse transesophageal irrigation of the mediastinum was achieved with orally ingested fluid. If the patient could not drink, the mediastinal irrigation was accomplished with a nasogastric tube positioned in the upper esophagus proximal to the perforation and connected to a bottle containing saline solution that was allowed to drip at a rate sufficient to keep the mediastinum clean. More precise guidelines were not available either in the original article¹ or in Dr. Santos' letter to the Editor.

Santos and Frater described eight patients who had a perforation not amenable to repair or a leak resulting from disruption of an attempted esophageal repair. It would be interesting to know how many primary repairs or other techniques were used during this time in their institution or whether these eight patients comprise their total experience with esophageal rupture. A critical examination of the series of patients presented by Santos and Frater¹ reveals principal differences compared with our patients.² Three of the eight patients had cervical esophageal rupture, which is a totally different condition from delayed septic perforation of the thoracic esophagus and has a very good prognosis even in delayed cases. The remaining five patients had rupture of the thoracic esophagus. All of our 34 patients² had a perforation of the thoracic esophagus with mediastinal sepsis, and the mean delay in diagnosis was 4 days, whereas the patients of Santos and Frater were operated on 26, 30, and 36 hours after rupture. This means that their diagnoses were made earlier and the delay in diagnosis may have been less than 1 day. The longer the diagnosis and treatment are delayed, the greater the mortality.³

Although the time factor and anatomic location are of critical importance in the prognosis of esophageal perforation, the size of the rupture also has an impact on the outcome. The size of the esophageal perforations in the report of Santos and Frater varied between 2.5 and 5 cm, whereas some of our patients had

tears measuring 10 cm.⁴ Taken all together, it is evident that the two series of patients are not comparable because of differences in the delay in diagnosis and the size of the perforations. Therefore no comparison of the reported therapeutic methods is possible.

I appreciate the interesting technique of transesophageal irrigation presented by Santos and Frater and hope that they report their results in delayed cases soon. However, I am dubious about its effectiveness. In delayed cases of mediastinitis the infection has spread far from the rupture in the mediastinum and pleural cavities. Open techniques (thoracotomy) allow exact placement of irrigation and drainage tubes, as well as careful cleansing of the mediastinum and both pleural cavities. Esophagectomy is necessary to remove the often gangrenous esophagus and to prevent the esophagomediastinal reflux of digestive gastric and duodenal juices, as well as intestinal bacteria. Cervical esophagostomy diverts the swallowed salivary enzymes and bacteria. We continue to use esophagectomy and mediastinal irrigation with local and intravenous antibiotics for the treatment of patients with delayed esophageal perforations with mediastinal sepsis. In early cases primary repair is the method of choice. If the perforation is located in the distal esophagus the primary repair site should be covered with gastric fundus and in more proximal perforations with omentum or pleura. Fibrin sealant (Tisseel, Immuno AG, Vienna, Austria) may be used for additional security.

*Jarmo A. Salo, MD
Department of Thoracic and Cardiovascular Surgery
Helsinki University Central Hospital
Helsinki, Finland*

REFERENCES

1. Santos GH, Frater WM. Transesophageal irrigation for the treatment of mediastinitis produced by esophageal rupture. *J THORAC CARDIOVASC SURG* 1986;91:57-62.
2. Salo JA, Isolauri JO, Heikkilä LJ, et al. Management of delayed esophageal perforation with mediastinal sepsis: Esophagectomy or primary repair? *J THORAC CARDIOVASC SURG* 1993;106:1088-91.
3. Orringer MB, Stirling MC. Esophagectomy for esophageal disruption. *Ann Thorac Surg* 1990;49:35-43.
4. Salo JA, Seppälä KMY, Pitkäranta PP, Kivilaakso EO. Spontaneous rupture and functional state of the esophagus. *Surgery* 1992;112:897-900.

Regression of autoimmune thrombocythemia after resection of pulmonary plasma cell granuloma

To the Editor:

Plasma cell granuloma (PCG), a rare, nonneoplastic lesion of the lung, first was described by Bahadori and Liebow.¹ PCG accounts for fewer than 1% of all lung tumors. It occurs at all ages but predominantly in children and young adults; sexes and ethnic groups are equally affected. Clinical features, including evidence of a preceding or concurrent respiratory infection, are variable, but almost 70% of patients have no symptoms.² Associations with such immunologic and hematologic disorders as hypergammaglobulinemia involving immunoglobulin G, M, and A fractions, elevation of erythrocyte sedimentation rate,