between benign and malignant adrenal masses. Burt and coworkers in fact refer to our 1992 article and acknowledge that chemical-shift MRI might be more effective than the methods they used. They did not, as they state, use the "best currently available MRI scanning techniques" for their 1994 publication.

I hope that Burt and associates and other thoracic and cardiovascular surgeons maintain their interest in the potential impact of MRI on presurgical diagnosis. Well-designed studies on this subject would be of benefit to the surgical and radiologic literature alike.

Donald G. Mitchell, MD
Professor of Radiology
Thomas Jefferson University Hospital
Jefferson Medical College
132 South 10th St.
196 Main Building
Philadelphia, PA 19107

REFERENCES

1. Burt M, Heelan RT, Coit D, et al. Prospective evaluation of unilateral adrenal masses in patients with operable non-small-cell lung cancer: impact of magnetic resonance imaging. *J THORAC CARDIOVASC SURG* 1994;107:584-9.
2. Kier R, McCarthy S. MR characterization of adrenal masses: field strength and pulse sequence considerations. *Radiology* 1989;171:671-4.
3. Baker ME, Blinder R, Spritzer C, Leight GS, Herfken RJ, Dunnick NR. MR evaluation of adrenal masses at 1.5 T. *AJR Am J Roentgenol* 1989;153:307-12.
4. Leroy-Willig A, Courtieu J, Roycayrol JC, Luton JP, Louvel A, Niesenbaum N. In vitro adrenal cortex lesions: characterization by NRM spectroscopy. *Magn Reson Imaging* 1987;5:339-44.
5. Leroy-Willig A, Bittoun J, Luton JP, et al. In vivo MR spectroscopic imaging of the adrenal glands: distinction between adenomas and carcinomas larger than 15 mm based on lipid content. *AJR Am J Roentgenol* 1989;153:771-3.
6. Mitchell DR, Crovello M, Matteucci T, Petersen RO, Miettinen MM. Benign adrenocortical masses: diagnosis with chemical shift MR imaging. *Radiology* 1992;185:345-51.
7. Tsushima Y, Ishizaka H, Matsumoto M. Adrenal masses: differentiation with chemical shift, fast low-angle shot MR imaging. *Radiology* 1993;186:705-9.

12/8/58953

Reply to the Editor:

In response to Dr. Donald G. Mitchell's letter to the Editor, we have the following comments. We thank Dr. Mitchell for his comments and criticisms. Although magnetic resonance imaging (MRI) methods are currently being investigated that possibly could lead to better discrimination of adrenal masses as being benign or metastatic in non-small-cell lung cancer, Dr. Mitchell did not take into account the time it takes for a scientific study to come to full publication. The inception of the study was

in February of 1990, and it was approved by our institutional review board in April of 1990. The accrual of the patients reported in our study began in August of 1990 and was completed in October of 1992. An abstract of our study was submitted to The American Association for Thoracic Surgery on November 11, 1993, and the paper was presented at the Seventy-third Annual Meeting in April of 1993. A manuscript was submitted at this meeting. This manuscript was accepted August 20, 1993, and published in the February 1994 issue of THE JOURNAL OF THORACIC AND CARDIOVASCULAR SURGERY. Dr. Mitchell published his technique of chemical shift MRI in the November 1992 issue of *Radiology*, after the completion of our study.

Although newer techniques are being developed in MRI to differentiate benign from malignant adrenal masses, few studies have histologic documentation of all patients reported. Therefore, although our study used the MRI techniques available during the period of the study, the point is still valid that the gold standard for determining the cause of adrenal masses is by cytopathologic or histopathologic study. We too are currently evaluating chemical-shift MRI and other newer MRI techniques, but we remain convinced that histologic examination is the gold standard by which to compare any new MRI technique.

Michael Burt, MD, PhD^a
Robert Heelan, MD^b
Department of Surgery^a
Department of Radiology^b
Memorial Sloan-Kettering Cancer Center
1275 York Ave.
New York, NY 10021
12/8/60339

Airway complications after lung transplantation: Is there a left-sided predilection?

To the Editor:

We read with interest the article by Griffith and associates¹ that was published in this JOURNAL. They have clearly shown that most complications of airway anastomoses after lung transplantation can be avoided by the refinement of surgical techniques (Table I, *J THORAC CARDIOVASC SURG* 1994;107:746). Although the etiology of impaired bronchial healing after lung transplantation is multifactorial, bronchial ischemia seems to be the primary cause.² In this series all seven patients with severe bronchial stenoses after bilateral lung transplantation had stenotic lesions of the left bronchus (three had bilateral stenoses). These patients were treated with endobronchial stents. Two patients had bronchial stenosis necessitating stenting after single left lung transplantations; by contrast, no bronchial stenoses occurred after single right lung transplantation (Table II, *J THORAC CARDIOVASC SURG* 1994;107:746). Colt and associates³ also have reported that all ten patients in whom stenosis developed after double lung transplantation demonstrated abnormalities at the level of the left main bronchial anastomosis (three had bilateral stenoses).