the first-generation Russian myocardiostimulators (Stim-nak 805, Moscow, Russia). Since 1991 the second-generation device (EKS 445, Moscow, Russia) has been used. The advantage of these stimulators is the ability to change stimulation ratios from 1:1 to 1:8. Four patients died during the early postoperative period, within 7 days after the operation (coronary artery thromboembolism in 1; progressive heart failure in 2; acute myocardial infarction in 1). One patient died in the postoperative period at 25 days; cause of death was cerebrovascular stroke. In the late postoperative period (8 and 12 months after operation) two patients have died of myocardial infarction. The follow-up period ranged from 6 months to 4.5 years. Fourteen patients were studied. Tolerance to physical exercise increased by two times up to 83.7 watts. The left ventricular end-diastolic volume decreased by an average of 15% to 265.1 ± 3.8 ml. The left ventricular ejection fraction increased by an average of 10% up to $41.2 \pm 1.6\%$. Ten patients were examined and selected to determine the optimal regimen of myostimulation. Mean value of the cardiac index without stimulation was 3.9 ± 1.2 L/min per square meter, and with stimulation it was 5.1 ± 1.2 L/min per square meter. Stroke index changed from 38.5 ± 3.2 ml/m² without stimulation to 51.0 ± 4.2 ml/m² with stimulation. The maximal increase of cardiac index and stroke index was seen at a low synchronization ratio (1:4 or 1:6).

Because of our findings, I believe that cardiomyoplasty leads to clinical improvement in patients who have been followed up for 5 years. Cardiomyoplasty has a positive effect on central hemodynamics. The myostimulation regimens of 1:4 and 1:6 are the most advantageous for graft stimulation, and skeletal muscle rest is an important factor in overall outcome.

Valeri Chekanov, MD, PhD
Milwaukee Heart Project
Sinai Samaritan Medical Center
945 North 12th St.
Milwaukee, WI 53201-0342
12/8/54591

Atherosclerotic changes in the internal thoracic artery

To the Editor:

We have read the letter by van Son and Smedts,¹ in which they present histologic studies of the internal thoracic artery (ITA) and suggest that "it may be beneficial not to use the distal 10% to 20% segment of the ITA because of its potentially primarily muscular content and, hence, increased tendency toward intimal thickening." We would like to present our results, which are similar to theirs.

We have done histologic studies of the more distal part of left ITA in 237 patients with coronary disease. Atherosclerosis was observed in 20 (8.4%) patients. In four (1.6%), intimal thickening was 25% to 50%. In 40 (16.8%) patients, the thickness of the intima was less than 25% of the lumen, and other abnormalities were present in eight patients.

Because of these results and those presented by van Son and Smedts, we believe that the most proximal portion

possible of the ITA should be used for coronary anastomoses.

J. Rubio-Alvarez, PhD
J. Sierra-Quiroga, PhD
D. Duran-Muñoz, PhD
J. B. Bengochea, PhD
Department of Cardiac Surgery
University Hospital
Santiago de Compostela, Spain

REFERENCE

1. van Son JA, Smedts F. Bifurcated ("Y") internal thoracic-coronary artery grafts [Letter]. *J THORAC CARDIOVASC SURG* 1993;106:945-6.

12/8/56692

Imaging of traumatic rupture of the aorta

To the Editor:

In their letter titled "Transesophageal Echocardiography for the Diagnosis of Traumatic Injury of the Thoracic Aorta" (*J THORAC CARDIOVASC SURG* 1994;107:638-40), Hengster, Furtwängler, and Pernthaler allege that transesophageal echocardiography (TEE) "has never been reported to be used after accident in a traumatized patient." In March 1991, a publication from the Mayo Clinic reported a case of traumatic rupture of the aorta diagnosed by TEE.¹ In June of 1991, the Allegheny General Hospital group published a series of 58 patients with chest trauma evaluated over 10 years in which the modality was used, with traumatic rupture of the aorta being demonstrated in three patients.² In 1993, the University of Kentucky reported seven cases of traumatic rupture of the aorta among their 69 patients with trauma studied by this technique.³ In early 1993, two similar instances were reported from Paris, France.⁴ In 1991, we instituted a (blinded) study of TEE in patients with trauma and, since the reports of the experience of the Pittsburgh and Lexington teams, have used it routinely.

More important than the issue of priority are the conclusions, based on a single patient, that TEE is "highly sensitive and specific" and avoids "invasive and time-consuming investigations." In our experience with the modality (along with aortography) in more than 50 patients with suspected traumatic rupture of the aorta, TEE has been useful in the diagnosis of this injury. However, the lesion was missed by TEE in two patients in whom traumatic rupture of the aorta was demonstrated by aortogram and confirmed by operation. In both patients, the aortogram was performed first and had demonstrated the transection. Despite intense efforts by experienced observers, the injury could not be visualized by means of TEE.

Because of the reasons mentioned in the authors' letter, TEE is a useful modality in patients with trauma. It is an adjunct but cannot completely supplant aortography. Our early experiences with helical or spiral computed tomographic studies are impressive, but we must remember that computed tomography itself was originally considered highly sensitive until tragic experiences taught otherwise. A high index of suspicion in an experienced clinician