abdominal aortic counterpulsating device. *Trans Am Soc Artif Intern Organs* 1984;30:540-4.

5. Utoh J, Whalen RL, Wilkerson BR, Fukamachi K, Harasaki H. Chronic in vivo function of a new ventricular assist device: the extracorporeal pulsatile assist device (EPAD). *Int J Artif Organs* 1993;16:91-5.

12/8/75204

Reply to the Editor:

Dr. Utoh makes the point that "the concept of this type of device is not so new." The counterpulsation technique has been used in many different ways. The application techniques were greatly improved to be more effective and suitable for long-term assistance.¹ The new idea for the improvement in the effectiveness of the counterpulsation technique was the use of high stroke volume in our device. This concept was the object of the discussion by Dr. Kantrowitz that followed the first presentation.² The theoretical explanation for the increased salutary hemodynamic effects of a high stroke volume counterpulsation device is based on the Suga-Sagawa law.³ On the basis of this law, for a given cardiac situation, the lower the left ventricular end-systolic pressure the higher the left ventricular stroke volume. For a counterpulsation device, the

higher the stroke volume the lower the left ventricular end-systolic pressure; thus, the higher the left ventricular stroke volume the better the systemic perfusion. The optimal counterpulsation volume in comparison with the capacitance of the arterial systemic bed has not been thoroughly investigated.

John N. Nanas, MD, PhD
Spyridon D. Mouloupoulos, MD, PhD
University of Athens School of Medicine
"Alexandra" Hospital
Athens, Greece

REFERENCES

1. Nanas J, Mouloupoulos S. Counterpulsation: historical background, technical improvements, hemodynamic and metabolic effects. *Cardiology* 1994;84:156-67.
2. Nanas J, Olsen D, Hamanaka Y, Chiang B-Y, Nielsen S, Hughes S, et al. Experience with a valveless, implantable abdominal aortic counterpulsating device. *Trans Am Soc Artif Intern Organs* 1984;30:540-4.
3. Suga H, Sagawa K. Instantaneous pressure-volume relationships and their ratio in the excised supported canine left ventricle. *Circ Res* 1974;35:117-26.

12/8/75203