daughter cysts. Atypical segmentectomy of the left lower lobe was performed with the help of GIA stapler (Auto Suture Company Division, United States Surgical Corporation, Norwalk, Conn.). At the level of the diaphragmatic dome, the pleura was decorticated. There was a splenopleural fistula, which was enlarged. Splenic cysts were aspirated transdiaphragmatically, removing 1 L of fluid, and thereafter the cavities were washed with hypertonic saline solution. External marsupialization over two left-sided transabdominal tubes was performed, and two left-sided chest drainage tubes were placed.

Pathologic examination found the pleural fluid and the lung specimen to contain daughter cysts and scolices. Postoperative recovery was uneventful, and the tubes were removed when they no longer drained. Mebendazole (500 mg every 8 hours) was administered for a month. There has been no recurrence for more than a year.

Splenic HD may be primary (more frequent³) or secondary, related to ruptured intraabdominal cysts. In primary infestation, the parasite has passed through hepatic and pulmonary filters.

Splenothoracic hydatidosis is a rare occurrence, usually occurring as a result of multivisceral intraabdominal hydatidosis.³ Recently, von Sinner and colleagues⁴ described a man with secondary intrathoracic hydatidosis. Previously, the patient had had diffuse intraabdominal HD (involving the spleen). Unlike that patient, our patient had solitary splenic hydatidosis and had shown no clinical evidence of disease.

Splenic hydatid cysts may produce serious complications, with rare reports of rupture into the pleura² and

into the bronchial system.² Because the hydatid fluid is highly antigenic, rupture into the pleural cavity results in chest pain, cough, implantation of the cyst in the cavity, and sometimes anaphylactic shock.

In our case, *Echinococcus* affected the thorax by extension of splenic HD across the diaphragm. Most likely, the growing cyst had previously eroded the pleura, resulting in the formation of a splenopleural fistula and later parenchymal involvement. There was a pleural effusion as well as multiple pleural hydatidosis caused by the development of the fistula.

Surgery remains the treatment of choice for HD. In associated hydatidosis of the thorax and abdomen, the operation must be carried out with thoracoabdominal access. When there are gross adhesions between the diaphragm and the liver (or spleen), however, it may be wise to primarily address surgical treatment of the chest lesion⁵ with abdominal cyst evacuation.

REFERENCES

1. Al-Mohaya S, Al-Awami M, Vaidya MP, Knox-Macaulay H. Hydatid cyst of the spleen. *Am J Trop Med Hyg* 1986;35:995-9.
2. Ríos Mozo M, Fernández Sanz S. Hidatidosis esplénica. *Rev Clin Esp* 1958;60:67-72.
3. Muñoz Sánchez JA, Conthe Gutiérrez P, Arnalich Fernández F, et al. Hidatidosis en un hospital general. I. Análisis epidemiológico de 1056 casos. *Med Clin* 1982;78:421-6.
4. Von Sinner WN, Rifai A, te Strake L, Sieck J. Magnetic resonance imaging of thoracic hydatid disease. *Acta Radiol* 1990;31:59-62.
5. Borrs S, Shaw J. Hepato-bronchial fistula caused by hydatid disease. *Thorax* 1981;35:25-8.

HEMOPTYSIS CAUSED BY AN ENDOBRONCHIAL LIPOMA

Antonio Piñero, MD, Alberto Giménez, MD, Fernando G. Lax, MD, and Pascual Parrilla, Murcia, Spain

Benign tumors of the lung and endobronchial tree are uncommon processes that, because of their location, need to be differentiated from malignant tumors. When these tumors are endobronchial, bronchoscopy is an excellent method of diagnosis. We present a case in which an endobronchial lipoma in a patient with symptoms of coughing and hemoptysis was removed by rigid bronchoscopy.

A 68-year-old male patient who had no family medical

history of interest, smoked approximately 20 cigarettes daily, and had a peptic ulcer was admitted to our hospital with a cough of several weeks' duration, accompanied by expectoration and episodes of hemoptysis. No other symptoms were reported. Neither general examination nor cardiopulmonary auscultation revealed pathologic findings.

Simple chest radiography disclosed condensation and atelectasis of the middle right lobe. The chest computed tomographic scan showed no mediastinal adenopathies. At bronchoscopy, a smooth, round, polypoid tumor, which was freely movable, apparently benign, and impossible to sample for biopsy, was seen within the lumen of the middle right lobe bronchus. Cytologic study from brushing revealed no neoplastic cells.

We subsequently performed a rigid bronchoscopy with the patient under general anesthesia. We took an

From the Department of General Surgery, Virgen de la Arrixaca, University Hospital, University of Murcia, Murcia, Spain.

Received for publication May 8, 1995; accepted for publication May 18, 1995.

J Thorac Cardiovasc Surg 1996;111:1104-5

Copyright © 1996 by Mosby-Year Book, Inc.

0022-5223/96 \$5.00 + 0 12/54/66328

Fig. 1. Nodule of mature adipose tissue enveloped in ciliate respiratory epithelium and separated by a fibrous area with inflammatory infiltrate. (hematoxylin-eosin stain; original magnification $\times 40$.)

initial biopsy sample and completely removed the tumor, which was pedunculated. Histologic examination of the resected piece showed a well-delimited, 7 mm diameter nodular proliferation. Although not clearly encapsulated, it was made up of mature adipose tissue with normal morphologic characteristics. The nodule was completely covered with ciliate respiratory epithelium of normal characteristics, nonulcerated and separated from proliferative adipocytes by a fibrous area with a slight subepithelial inflammatory infiltrate. Diagnosis of endobronchial lipoma was made (Fig. 1).

The patient was discharged 48 hours after operation. He has been free of symptoms at subsequent follow-up examinations.

Benign pulmonary tumors are rare entities, and among them lipomas are the most uncommon. Of a total of 3502 pulmonary tumors, Jensen and Petersen¹ reported only 65 cases of benign pulmonary tumors, only three of which were lipomas. The medical literature in the English language as of 1991 had reported only 54 cases of endobronchial and endotracheal lipoma.

Classically, endothoracic lipomas have been divided into endobronchial, parenchymatous, pleural, mediastinal, and cardiac.³ The endobronchial type originates in the submucous adipose tissue of the bronchial tree, grows endoluminally, and usually manifests as a freely movable, pedunculated tumor with a narrow stem.

The clinical symptoms of endobronchial lipoma are fundamentally caused by obstruction of the airflow, with

the possibility of distal atelectasis, overinfections, and pneumonia. Although coughing has been reported as a clinical manifestation in 81%, hemoptysis was present in only 26%.³ Differential diagnosis must be made with carcinoma and adenoma. Bronchoscopic diagnosis to differentiate endobronchial lipomas from adenomas can be difficult. The surface of the tumor, however, is paler in the lipoma and more friable in the adenoma.⁴

These tumors can be managed by bronchoscopy, as in the case reported here, when practicable and the peduncle permits. If bronchoscopic management is not possible, we must resort to thoracotomy with bronchiotomy and removal of the tumor before the symptoms have led to irreversible distal complications.

REFERENCES

1. Jensen MS, Petersen AH. Bronchial lipoma: three cases and review of the literature. *Scand J Thorac Cardiovasc Surg* 1970;4:131-4.
2. Matsuba K, Saito T, Ando K, Shirakusa T. Atypical lipoma of the lung. *Thorax* 1991;46:685.
3. Politis J, Funahashi A, Gehlsen JA, DeCock D, Stengel BF, Choi H. Intrathoracic lipomas: report of three cases and review of the literature with emphasis on endobronchial lipoma. *J Thorac Cardiovasc Surg* 1979;77:550-6.
4. Bellin HJ, Libshitz HI, Patchefsky AS. Bronchial lipoma: report of two cases showing chondroitic metaplasia. *Arch Pathol* 1971;92:20-3.