LETTERS TO THE EDITOR

Partial sternotomy: A mini approach for maxi exposure

To the Editor:

In response to a letter to the Editor by Moreno-Cabral in the Journal (1997;113:810-1), I want to endorse his enthusiasm for the partial lower sternotomy. I, too, have been frustrated with the limitations of exposure of the heart via left and right anterolateral thoracotomy. In contrast, partial lower sternotomy up to the second intercostal space with bilateral transection of the sternum at the top of the incision allows adequate exposure for a wide variety of operations.

Over the past 3 months, I have performed 29 operations using partial lower sternotomy. All patients received from one to five coronary bypasses; in 26, the left internal thoracic artery was used and, in one, the right internal thoracic artery. In addition, two had resection of a left ventricular aneurysm, one replacement of both the aortic and mitral valves, and two replacement of the aortic valve. In one patient (coronary bypass plus double valve replacement), the sternal incision was extended to the first intercostal space and the sternum was transected at that level.

Postoperative pain was less than in patient with anterolateral thoracotomies. Blood loss was less than in patients with complete sternotomy. The skin incision was shorter than with complete sternotomy and only slightly longer than that need for the anterolateral thoracotomy approach.

Partial sternotomy is a worthwhile alternative to complete sternotomy and has many of the advantages of other minimally invasive techniques.

Hans Zwart, MD
Suite 322
580 Lincoln Park Blvd.
Kettering, OH 45429-3493
12/8/83790

The right internal thoracic artery for myocardial revascularization

To the Editor:

We read with great interest the report by Ueyama and associates (J Thorac Cardiovasc Surg 1996;112:731-6) regarding revascularization of the posterolateral myocardial wall with the right internal thoracic artery (ITA) graft via the transverse sinus. This technique, published by us¹ in 1984 as a complementary method to use with the left ITA to the left anterior descending branch, provoked several comments concerning early morbidity, such as hemorrhage, mediastinitis, and late occlusion of the graft. On this subject, publications by Galbut,² Buche,³ and Ueyama and colleagues,⁴ as well as our own report,⁵ show that the incidence of reoperation for bleeding ranged from 1.9% to 2.6% and mediastinitis from 0.9% to 2.6%, values normally found with any cardiac operation. The

late patency rate of the right ITA through the transverse sinus and grafted to the circumflex artery or obtuse marginal branch ranged from 80.6% to 98%.^{2,3,5} In our 233 patients who received both ITAs with or without saphenous vein grafts or the inferior epigastric artery, there were 49 late recatheterizations (average 61.9 months) and the patency of the grafts was as follows: 45 (92%) right ITA grafts, 47 (96%) left ITA grafts, and 19 (70%) saphenous vein grafts. In symptomatic patients with both ITAs patent, reoperation was never necessary. Only one patient required reoperation, 6 months after the original operation, because of the occlusion of the two ITAs. The great advantage of using ITAs or other arterial grafts for the left anterior descending branch and the obtuse marginal branch of the circumflex artery is to avoid late reoperation. We think that in due time we will choose the arterial graft most appropriate for each specific coronary artery. On the basis of our results, we continue to choose both ITAs for revascularization of the branches of the left coronary artery in patients up to 70 years of age who do not have excessive obesity, severe diabetes, or other important risk factors.

Luiz B. Puig, MD
University of São Paulo Medical School
Cardiopulmonary Department
São Paulo, Brazil

REFERENCES

1. Puig LB, Neto LF, Rati M, Ramires JAF, Luz PL, Pileggi F, et al. A technique of anastomosis of the right internal mammary artery to the circumflex artery and its branches. *Ann Thorac Surg* 1984;38:533-4.
2. Galbut DL, Traad EA, Dorman MJ, DeVitt PL, Larsen PB, Kurlansky PA, et al. Coronary bypass grafting in the elderly. *J Thorac Cardiovasc Surg* 1993;106:128-36.
3. Buche M, Schroeder E, Chenu P, Gurne O, Marchandise B, Pompilio G, et al. Revascularization of the circumflex artery with the pedicled right internal thoracic artery: clinical, functional, and angiographic midterm results. *J Thorac Cardiovasc Surg* 1995;110:1338-43.
4. Ueyama K, Sakata R, Umebayashi Y, Nakayama Y, Arakari K, Ura M. In situ right internal thoracic artery graft via transverse sinus for revascularization of posterolateral wall: early results in 116 cases. *J Thorac Cardiovasc Surg* 1996;112:731-6.
5. Gerola R, Puig LB, Moreira LFP, Cividanes GV, Gemha GP, Souto RCM, et al. Right internal thoracic artery through the transverse sinus in myocardial revascularization. *Ann Thorac Surg* 1996;61:1708-13.

12/8/79882

Reply to the Editor:

We were intrigued with Dr. Puig's response. We also believe that use of the right internal thoracic artery (ITA) via the transverse sinus to revascularize the circumflex

area in coronary artery bypass operations is theoretically the best procedure.

Regarding indications for use of bilateral ITAs, our opinion on age limit differs from Dr. Puig's. Our age limit for bilateral ITAs is 75 years in Japan, because the average life span in Japan is 80 years. We assume that it is shorter in Brazil.

Since publishing our report in the Journal, we have continued to use this technique for selected patients. The total number of patients so treated as of November 28, 1996, is 215. Excellent long-term patency of the right ITA grafts via the transverse sinus in Dr. Puig's experience reinforces our results and gives us confidence in using this method.

*Koji Ueyama, MD
Ryuzo Sakata, MD
Department of Cardiovascular Surgery
Kumamoto Central Hospital
1-16-1 Shinyashiki
Kumamoto 862, Japan
12/8/79881*

Early tracheal extubation after coronary artery bypass grafting

To the Editor:

We read with great interest the report by Cheng and colleagues¹ about early versus conventional tracheal extubation after coronary artery bypass grafting. This subject is of great importance in areas in which resources are limited.

In July 1976 we² published our first article on early extubation in general cardiac surgery. In support of early extubation after valve replacement, Midell and associates³ in 1974 published their experience with 100 cases, and Barash and coworkers⁴ of Yale University, in 1980, showed that only 4% of patients undergoing pediatric cardiac operations required reintubation. In August of 1980, in a thesis to become professor of surgery at the University of Ceará Medical School, Jucá⁵ reported the results in 105 patients undergoing coronary bypass grafting who were subjected to early extubation and 20 patients who were extubated in the operating room. Nowadays, if patients have had mild body hypothermia (30° to 34° C) during perfusion, we prefer to extubate them 1 to 4 hours after arrival in the postoperative intensive care unit. The beneficial effects of early extubation on ventricular performance were demonstrated in 1988 by the group from Duke University.⁶

We appreciate the work of the group from the Toronto Hospital because their subject is of great importance in undeveloped areas like ours in northeast Brazil.

*Regis Jucá, MD, FACS
Professor of Surgery
University of Ceará
Lery Monte
Anesthesiology
Casa de Saúde São Raimundo
Ceará, Brazil*

REFERENCES

1. Cheng DCH, Karski J, Peniston C, et al. Morbidity outcome in early versus conventional tracheal extubation after coronary artery bypass grafting: a prospective randomized controlled trial. *J Thorac Cardiovasc Surg* 1996;112:775.
2. Jucá ER, Rolim JM, Monte L, et al. Extubação precoce após cirurgia cardíaca com C.E.C.. *Rev Col Bras Cirurg* 1976;julho-agosto;163.
3. Midell AI, Skinner DB, De Buer A, Bermudez G. Review of pulmonary problems following valve replacement in 100 conservative patients: case against routine use of assisted ventilation. *Ann Thorac Surg* 1974;18:278.
4. Barash PG, Lescovich F, Katz J, Falner WS, Stansel HC. Early extubation following pediatric cardiothoracic operation: a viable alternative. *Ann Thorac Surg* 1980;29:228.
5. Jucá ER. Extubação traqueal precoce após cirurgia de revascularização do miocárdio com enxerto de veia safena aorto-coronária [thesis]. Ceará, Brazil: University of Ceará; 1980.
6. Gall J Jr, Olsen CD, Reves JC, et al. Beneficial effects of endotracheal extubation on ventricular performance. *J Thorac Cardiovasc Surg* 1988;95:819-27.

12/8/83786

Reply to the Editor:

I would like to thank Jucá and Monte for their interest and support of our prospective, randomized, controlled trial on morbidity outcome in early versus conventional tracheal extubation after coronary artery bypass graft operations.¹ As correctly indicated in their letter, the practice of early tracheal extubation in cardiac surgery is even more important in places with limited resources in this time of cost containment and environment. As a matter of fact, we² have recently published a comprehensive cost study of early tracheal extubation in patients undergoing coronary bypass, including preoperative heart catheterization costs, physician fees, and all departmental costs from patient admission to discharge from the hospital. This study demonstrated that early tracheal extubation significantly reduces the cost of coronary artery bypass grafting by reducing the variable costs in the intensive care unit, reducing the length of stay in the intensive care unit, and reducing the postoperative hospital stay. This cost study further emphasizes that early extubation in these patients does not increase the postoperative morbidity or the cost of postoperative complications, as major postoperative complications were indeed very costly as documented.² On the subject of resource use, early tracheal extubation resulted in fewer cancellations of cardiac operations by improving the use of the intensive care unit and allowed an increase in the caseload of cardiac surgery.

I congratulate Jucá and colleagues³ on their earlier work and insight on early extubation in cardiac surgery. As indicated in our morbidity article,¹ not only has the routine use of prolonged controlled ventilation in patients undergoing cardiac operations been standard practice for the past three decades, but early tracheal extubation in these patients is not new. Cost containment and efficient resource use force the pendulum back to the debate of early tracheal extubation in patients undergoing cardiac operations. As we indicated in our morbidity article,¹ most of these earlier studies were observational studies, case reports, and retrospective reviews with minimal outcome measurements. The most common