a protocol of using the technique for established cases of endocarditis, in which the incidence of prosthetic valve endocarditis is highest.

It is logical to consider the concept of impregnating all valves with antibiotics to minimize prosthetic valve endocarditis caused by perioperative contamination. We suggest that to minimize crossclamp time but to provide adequate short-term local antibiotic levels in the valve sewing ring, a commercially available valve with a protein-sealed sewing ring could be soaked with rifampicin, similar to established techniques used in vascular surgery.^{2,3}

*B. G. French, FRACS
M. F. O'Brien, FRACS
Department of Cardiac Surgery
The Prince Charles Hospital
Brisbane 4032, Australia*

REFERENCES

1. Karck M, Siclari F, Wahlig H, Sperling U, Schmid C, Haverich A. Pretreatment of prosthetic valve sewing ring with antibiotic fibrin sealant compound as a prophylactic tool against prosthetic valve endocarditis. *Eur J Cardiothorac Surg* 1990;4:142-6.
2. Strachan CJL, Newsom SWB, Ashton TR. The clinical use of an antibiotic-bonded graft. *Eur J Vasc Surg* 1991;5:627-32.
3. Avramovic J, Fletcher JP. Prevention of prosthetic vascular graft infection by rifampicin impregnation of a protein sealed Dacron graft in combination with parenteral cephalosporin. *J Cardiovasc Surg* 1992;33:70-4.

12/8/81586

Measurement of aortic diameter

To the Editor:

In the October 1996 issue of the Journal, Weinert and colleagues¹ give an elegant account of their measurements of aortic diameter by means of multiplane transesophageal echocardiography. I have no complaints whatsoever concerning the excellence of the techniques described, but I have major problems with the anatomic structures that are purported to be measured. Apart from the fact that there is no ringlike structure^{2,3} supporting the hinge-points of the aortic valve leaflets (the so-called "anulus"), if the Chicago group makes measurements as they claim, then it is impossible for them to quantify the fullest diameter of the aortic root. The reason, of course, is that the aortic valve possesses three sinuses and leaflets and, even though the sinuses are rarely completely equal,⁴ measuring from the basal attachment of the noncoronary leaflet to the basal attachment of the right coronary leaflet will always produce a chord of the outflow tract. Calculations on the basis of the aortic root being a circle shows that measurements of the kind proposed underestimate the true diameter by a factor of 10% or more. The aortic root, of course, is not a true circle, but these geometric

considerations are surely worthy of at least some discussion in studies of this kind.

*Robert H. Anderson, MD, FRCPath
Imperial College School of Medicine
Department of Paediatrics
National Heart and Lung Institute
Dovehouse St.
London SW3 6LY, United Kingdom*

REFERENCES

1. Weinert L, Karp R, Vignon P, Bales A, Lang RM. Feasibility of aortic diameter measurement by multiplane transesophageal echocardiography for preoperative selection and preparation of homograft aortic valves. *J Thorac Cardiovasc Surg* 1996;112:954-61.
2. Anderson RH, Devine WA, Ho SY, Smith A, McKay R. The myth of the aortic annulus: the anatomy of the subaortic outflow tract. *Ann Thorac Surg* 1991;52:640-6.
3. Sutton JP III, Ho SY, Anderson RH. The forgotten interleaflet triangles: a review of the surgical anatomy of the aortic valve. *Ann Thorac Surg* 1995;59:419-27.
4. Vollebergh FEMG, Becker AE. Minor congenital variations of cusp size in tricuspid aortic valves: possible link with aortic stenosis. *Br Heart J* 1977;39:1006-11.

[Response declined]

12/8/81128

Neuropsychologic outcome after normothermic cardiopulmonary bypass

To the Editor:

We agree happily with virtually all of Professor Treasure's commentary following our paper.¹ Indeed, our agreement is already explicit in the text. We planned and executed a randomized controlled study within pre-recognized externally applied constraints on numbers. We considered the available outcome measures and concluded that none would provide sufficient power to detect differences of any interesting size with the numbers available. We welcome the consensus of experts that has taken place since our study was planned and completed, but know that the consensus is as much a way of sharing responsibility for error as a guaranteed pathway to truth. With the information available at the time, we made a prospective choice of an outcome measure that should logically have been more sensitive than existing ones and declared it to be our primary measure. No pilot data were available on this measure that could have informed prospective power calculations. It proved, in any case, to have enough power to detect an interesting difference with 95% confidence, although we made no secret of our doubt as to the clinical meaning of the difference. We identified the further analysis as being subsidiary, to inform further studies on the problems of this nature.

Nowhere did we use our results to draw inferences about stroke rate. Nowhere did we draw any conclusions about safety. Our concluding sentence remarked that our results, and the absence of any separate demonstration of benefit, provided little incentive to recommend the routine use of normothermic systemic perfusion (instead of