aprotinin package insert provided by the drug's manufacturer.

The well-designed, randomized, blinded, placebo-controlled study reported by Vander Salm and associates is of value to cardiac surgeons and anesthesiologists in that it provides data regarding the efficacy (or the lack thereof) of EACA. As a surgeon who administers EACA on a regular basis and aprotinin to selected patients, this information is of interest to me. In my opinion, however, the authors' conclusions regarding the relative efficacy of EACA and their comparisons with aprotinin do not fit the results presented and should therefore be carefully considered by the reader.

*John H. Lemmer, Jr., MD
Northwest Surgical Associates
2226 N.W. Pettygrove
Portland, OR 97210-2608*

REFERENCES

1. Vander Salm TJ, Kaur S, Lancey RA, Okike ON, Pezzella AT, Stahl RF, et al. Reduction of bleeding after heart operations through the prophylactic use of epsilon-aminocaproic acid. *J Thorac Cardiovasc Surg* 1996;112:1098-107.
2. Davis R, Whittington R. Aprotinin: a review of its pharmacology and therapeutic efficacy in reducing blood loss associated with cardiac surgery. *Drugs* 1995;49:954-83.
3. Baele PL, Rui-Gomez J, Londot C, Sauvage M, Van Dyck MJ, Robert A. Systematic use of aprotinin in cardiac surgery: influence on total homologous exposure and hospital costs. *Acta Anesth Belg* 1992;43:103-12.
4. Lathi KG, Hariawala M, Fotouhi MD, Symes JF. Economics of aprotinin in cardiac surgery. *Anesth Analg* 1995;80:SCA119.
5. Van Norman G, Lu J, Spiess B, Soltow L, Gillies B. Aprotinin versus aminocaproic acid in moderate-to-high-risk cardiac surgery: relative efficacy and costs. *Anesth Analg* 1995;80:SCA19.
6. Laub GW, Riebman JB, Chen C, Adkins MS, Anderson WA, Fernandez J, et al. The impact of aprotinin on coronary artery bypass graft patency. *Chest* 1994;106:1370-5.
7. Lemmer JH Jr, Stanford W, Bonney SL, Breen JF, Chomka EV, Eldredge WJ, et al. Aprotinin for coronary bypass operations: efficacy, safety, and influence on early saphenous vein graft patency. A multicenter, randomized, double-blind, placebo-controlled study. *J Thorac Cardiovasc Surg* 1994;107:543-53.
8. Royston D, Taylor KM, Bidstrup BP, Sapsford RN. Effect of aprotinin on need for blood transfusion after repeat open heart surgery. *Lancet* 1987;2:1289-91.

12/8/80547

Reply to the Editor:

As Lemmer points out, correctly, our recent paper documented a reduction in chest tube drainage during the first 24 hours of about 200 ml after the use of epsilon-aminocaproic acid (EACA).¹ In that same series of patients, we did not observe a reduction in blood product administration.

We disagree, however, with his characterization of the reduction in blood as "essentially negative results." In

fact, the 200 ml incremental blood loss in the control group represents a 30% increase ($p = 0.002$). One reason for the difference of (only) about 200 ml between the control and EACA groups is that the patients operated on by surgeon 1 in the study had significantly less bleeding than the other patients, and surgeon 1 operated on a disproportionately large number of patients in the control group: 65% of the control patients were operated on by surgeon 1, and 56% of surgeon 1's patients were control patients. The second reason is that the overall blood loss in the series was low: 647 ml and 839 ml at 24 hours in the EACA and control groups, respectively. After reading Lemmer's criticisms, we almost feel obliged to apologize for our institution's good results. Had we a higher rate of blood loss, the difference between the control and EACA groups might have been greater, and a significant difference in blood product administration might also have been observed.

Except for the comparison of the cost of the two drugs, we did not compare EACA with aprotinin. Because Lemmer suggests that we drew conclusions regarding the relative efficacy of EACA and aprotinin (we did not), it may be of some use now to compare the results in our recent study with those of the excellent and conclusive multiinstitutional study of aprotinin reported by Lemmer and colleagues.² Our 24-hour cumulative blood losses were 647 ml in the EACA group and 839 ml in the control group. In Lemmer's report, the blood losses in primary coronary artery bypass graft (CABG) cohort at the time of chest tube removal were 855 ml and 1503 ml in the aprotinin and control groups, respectively. Inasmuch as the chest tubes in the Lemmer series were removed earlier (20.9 hours and 17.3 hours in the aprotinin and control groups, respectively) than in our patients (postoperative day 2), the hourly drainage rate in our series (calculated at 24 hours) is considerably less than in the Lemmer series (Table I). In their redo CABG cohort, the losses were 1225 ml and 1979 ml in the aprotinin and control groups, respectively. The results we reported are for all types of cardiac operations, including valve operations and redo operations. Thus, in this group of patients expected to be at higher risk of bleeding than were the patients reported on by Lemmer, our control patients had about as much bleeding as did the aprotinin-treated patients in Lemmer's series, and our treatment (with EACA) patients has less bleeding.

Although our red cell administration rate for the patients receiving EACA was slightly less than for the control patients, the difference was not statistically significant. Again, comparison with the patients in Lemmer's series serves to put the transfusion rates in proper context. Our transfusion threshold was a hematocrit value of 25%; in Lemmer's series it was 21%. With a higher hematocrit threshold in our series than in Lemmer's, and a study group that included complex valve operations as well as redo operations, a higher rate of transfusion would be expected in our series. Our EACA and control groups had a mean of 1.13 units and 1.35 units of packed red cells transfused, respectively; in Lemmer's series, the aprotinin and control groups had 1.1 and 2.1 units transfused, respectively. Our treatment and control groups had blood replacement at close to the level of Lemmer's aprotinin-

Table I

	<i>Vander Salm series (all patients)</i>		<i>Lemmer series (primary CABG)</i>	
	<i>Control</i>	<i>EACA</i>	<i>Control</i>	<i>Aprotinin</i>
Chest tube drainage rate (ml/hr)	35	27	87	41
Chest tube drainage (ml)	839	647	1503	855
Mean No. units of RBCs transfused	1.35	1.13	2.1	1.1

RBCs, Red blood cells.

treated group. In our series, the mean units of red cells transfused in all patients (both EACA and control groups) with primary operations was only 0.84, a number less than that of both the control and aprotinin-treated groups receiving a primary CABG in Lemmer's series. In a study of primary CABG operations reported by Daily and colleagues,³ the number of patients receiving transfusions was significantly less in the EACA-treated patients (4.8%) than in those in the control group (26.3%).

Lemmer points out that we did not compare aprotinin against EACA. He is correct. We did not intend to do so. We only wished to ask whether EACA reduces blood loss when compared with a control series not receiving EACA. The only comparison of EACA with aprotinin made in our paper was of the cost of the two drugs. We did not claim that EACA more effectively reduces bleeding than does aprotinin, only that it does so less expensively. In fact, we are currently engaged in a Bayer-supported study of aprotinin to ascertain whether its documented reduction of bleeding reduces transfusion costs and other intensive care unit costs (such as ventilator support) sufficiently to offset the high cost of the drug.

We are embarrassed to realize that our paper contains an egregious numerical error—as Lemmer noticed and we did not. In the original manuscript, the cost of aprotinin was based on the (correct) volume use of the high-dose regimen: 200 ml before institution of cardiopulmonary bypass, 200 ml in the heart-lung pump prime, and an infusion rate during the operation of 50 ml/hr. When the final manuscript was submitted, the aprotinin dose was converted from milliliters to kallikrein inactivator units (kIU) and a tenfold error was introduced. None of the authors noticed the error. The correct doses are 2×10^6 kIU before cardiopulmonary bypass, 2×10^6 kIU in the pump prime, and 0.5×10^6 kIU/hr infusion—not ten times that dose, as was published. However, since the cost calculation was based on the volume dose used, the cost calculation of \$785 per patient is correct as published, and the comparison with the \$6.96 per-patient EACA cost remains rather striking. We apologize for the error, which, fortunately, did not affect the cost calculation in our paper. When we use aprotinin, we use the standard high-dose regimen as described above and in the package insert.

Although Lemmer is critical of our conclusions, we suspect that we both have nearly identical indications for the use of aprotinin. As does he, we use it only in operations expected to carry a high bleeding risk. We probably differ in that we do not use the drug routinely in

redo operations because we do not consider those to carry a high bleeding risk.

Thomas J. Vander Salm, MD
Department of Surgery

University of Massachusetts Medical Center
55 Lake Ave., N.
Worcester, MA 01655-0304

REFERENCES

1. Vander Salm TJ, Kaur S, Lancey RA, Okike ON, Pezzella AT, Stahl RF, et al. Reduction of bleeding after heart operations through the prophylactic use of epsilon-aminocaproic acid. *J Thorac Cardiovasc Surg* 1996;112:1098-107.
2. Lemmer JH Jr, Stanford W, Bonney SL, Breen JF, Chomka EV, Eldredge WJ, et al. Aprotinin for coronary bypass operations: efficacy, safety, and influence on early saphenous vein graft patency. A multicenter, randomized, double-blind, placebo-controlled study. *J Thorac Cardiovasc Surg* 1994;107:543-53.
3. Daily PO, Lamphere JA, Dembitsky WP, Adamson RM, Dans NF. Effect of prophylactic epsilon-aminocaproic acid on blood loss and transfusion requirements in patients undergoing first-time coronary artery bypass grafting: a randomized, prospective, double-blind study. *J Thorac Cardiovasc Surg* 1994;108:99-106.

12/8/80545

Twenty-three-year survival of a bovine pericardial bioprosthesis

To the Editor:

The letter by Trujillo and colleagues¹ concerning a 19-year survival of a Hancock I porcine mitral bioprosthesis (Johnson & Johnson Cardiovascular, King of Prussia, Pa.) prompts us to report this 23-year survival of a bovine pericardial bioprosthesis in the aortic position.

A male patient was noted to have a cardiac murmur on routine medical examination at the age of 11 years. In 1972, at the age of 36 years, he was referred for treatment because of a 3-year history of dizziness, double vision, and progressive dyspnea on moderate exertion. A diagnosis of aortic stenosis was made. The patient was not evaluated by cardiac catheterization. On October 10, 1972, he underwent aortic valve replacement. A heavily calcified bicuspid aortic valve was excised and replaced with a 20 mm standard-profile Ionescu pericardial xenograft that had been constructed by the surgical team at the Leeds General Infirmary. He made an uncomplicated postoper-