LETTERS TO THE EDITOR

Predicting survivorship after coronary artery bypass grafting in patients with low ejection fraction

To the Editor:

We read with great interest an editorial by Dr. Wechsler¹ on coronary bypass grafting in patients with an ejection fraction of 20% or less. In this editorial, Dr. Wechsler emphasized the need to identify a subset of patients who benefit from coronary artery bypass grafting, regardless of the presence of low ejection fraction. He stated that a predictive scoring system was lacking that took into account most of the clinical and hemodynamic parameters as descriptors of left ventricular performance. He also claimed that he had not seen a series that combined parameters mathematically to evolve a risk score based on the important interaction between the critical risk elements.

In this letter, we would like to point out our recent study² published in the Journal in 1995, in which we tried to develop exactly the indicator Dr. Wechsler describes. In our study we sought to elucidate the relation between the preoperative parameters and the survivorship after coronary artery bypass grafting for patients with low ejection fraction (mean ejection fraction of $23\% \pm 7\%$). We used a technique called *factor analysis* to "summarize" a number of preoperative parameters. Preoperative clinical variables, as well as hemodynamic parameters from exercise stress and nitroglycerin challenge tests, were screened and entered into factor analysis to produce a score for predicting the life status (alive or dead) 4 years after the operation. We found that the most highly loaded parameters predicting survivorship were related to the exercise stress test (specifically, cardiac index and systemic vascular resistance) and nitroglycerin challenge (specifically, stroke index and cardiac index). A history of congestive heart failure was also predictive. Ejection fraction itself and the response of ejection fraction to exercise stress or nitroglycerin challenge were not useful parameters to predict the outcome. More important, a newly developed score (which we called *cardiac function factor*), derived from factor analysis, provided a predictive value superior to that of any individual variable. By performing logistic regression analysis, we were able to predict survival by entering individual variables to calculate cardiac function factor.

Our series was relatively small (56 patients), and the study was not done prospectively. For surgeons and cardiologists interested, prospectively, in studying patients with low ejection fraction, we strongly encourage them to include a preoperative exercise stress test in the protocol. In our study, the 7-year survival was 78% in the patients who had a cardiac index greater than 3.5 L/min per square meter at exercise, whereas only 29% of patients survived who had an exercise cardiac index less than 3.5 L/min per square meter. This is an easy number to obtain. Maximal oxygen consumption may be used as a noninvasive alternative to cardiac index. Left ventricular volume or echocardiographically measured ventricular dimension should

be determined. The usual formula of "large ventricle = bad ventricle" may not apply in this subset of patients.³ If possible, a load-insensitive parameter of contractility and the diastolic property of the ventricle should be taken into account. The underlying mechanism of altered survivorship should also be sought. We agree with Dr. Wechsler that much remains much to be done to identify the subset of patients with low ejection fraction who benefit from coronary artery bypass grafting. As Dr. Wechsler pointed out, this is an area still ripe for new investigation.

Shigeki Morita, MD
Guo-Wen Sun, PhD
Gregory Louis Kay, MD
The Heart Institute
Good Samaritan Hospital
123 S. Alvarado St.
Los Angeles, CA 90057

REFERENCES

1. Wechsler AS. Editorial on "Coronary artery bypass grafting in patients with an ejection fraction of twenty percent or less." *J Thorac Cardiovasc Surg* 1996;111:998-1000.
2. Morita S, Zubiato P, Kay GL, et al. Prediction of long-term survival by preoperative exercise testing in patients with depressed ejection fraction undergoing myocardial revascularization. *J Thorac Cardiovasc Surg* 1995;110:944-51.
3. Morita S, Kay GL, Zubiato P, Mendez M, Kay JH. Diastolic function determines the cardiac output in patients with low ejection fraction secondary to ischemic heart disease. *Circulation* 1989;80(Suppl):II275.

12/8/78758

Reply to the Editor:

The response to my editorial by Dr. Morita and his colleagues is much appreciated. I was aware of their article before the time of my writing. They are to be commended for taking on a tough task and using a sound method for establishing a predictive index of long-term survival after coronary bypass grafting in patients with reduced ejection fraction. The concept of factor analysis is sound and can serve as an amplifier of parameters that are independently of marginal significance but which, when combined, become more meaningful. Their article had a few shortcomings, which they acknowledged. These include the retrospective nature of their study, small numbers that may have been inadequately powered to detect important differences between groups, and an absence of focus on perioperative mortality. The wisdom of their approach was in combining an exercise variable that served as a surrogate for the efficacy of adaptive mechanisms in compensating for diminished ejection fraction in their patient population. As regards predicting operative mortality, similar measured parameters should be helpful but might be better analyzed in cohorts of patients in which the focus is even more heavily on those with low ejection fractions. In addition, readers of all such articles are reminded that every patient entered into the

series passed an initial screening test that suggested to the surgeon the likelihood of a good outcome from operative intervention.

Andrew S. Wechsler, MD
Medical College of Virginia
P.O. Box 980645
Richmond, VA 23298-0645
12/8/78747

Atrioventricular septal defect with subaortic stenosis: Extended valvular detachment and leaflet augmentation

To the Editor:

We read with interest the recent report by Van Ardsell and colleagues¹ regarding the different treatment options for subaortic stenosis associated with atrioventricular septal defect (AVSD). They pointed out the importance of leaflet augmentation and fibromyectomy to decrease the likelihood of recurrence of subaortic stenosis. We would like to further clarify the technical aspects of leaflet augmentation in AVSD when malattachment of the left superior leaflet is the predominant feature of an associated subaortic stenosis. The authors, when describing this surgical procedure, stated that "the superior bridging leaflet is detached from the anterior portion of the septal crest extending to the region just to the left of the left ventricular outflow tract" and that "the increased mitral aortic separation is excised." The left superior leaflet is thus detached at two different boundaries of the left ventricular outflow tract: (1) the septal insertion of the valvular leaflet and (2) the aorta-left atrioventricular (AV) valve continuity.

Leaflet augmentation with detachment of the valvular leaflet from the septal crest was first reported by Lappen and

colleagues² and subsequently by DeLeon and colleagues³ in a patient in whom subaortic stenosis recurred after the surgical correction of a partial AVSD. The anatomic concepts leading to that treatment option were emphasized by Chang and Becker⁴ in Rastelli type A AVSD. Today, this procedure gains increased interest⁵ owing to the failure of more simple options that did not address the pathoanatomic features of the subaortic area in AVSD.

The importance of aorta-left AV valve continuity in the pathogenesis of a subaortic stenosis was underlined by Ebels and coworkers⁶ in their description of the "atrial fold." These authors proposed a sham repair with either an enlargement of this anatomic region using a diamond-shaped patch or a resection of the atrial fold in case of an associated left AV valve replacement. Moreover, patch insertion at this level associated with a left AV valve replacement avoids postoperative subaortic stenosis in AVSD.⁷

Keeping these important anatomic concepts in mind, we used leaflet augmentation in a 12-year-old girl who had been referred to our institution for native subaortic stenosis associated with an AVSD. Leaflet augmentation was accomplished by extending the detachment of the superior leaflet from the septal crest into the aorta-left AV valve continuity (Fig. 1).⁸ The young girl was still doing well 2 years later, without subaortic stenosis, with a moderate stenosis at the aortic level and a mild "mitral" insufficiency. Since this report, we observed, immediately after correction of a partial AVSD in a 3-year-old girl without Down syndrome, the occurrence of a subaortic stenosis with a systolic left ventricle-ascending aorta gradient of 37 mm Hg. One year later, a color-coded echocardiographic study revealed the progressive increase of the left ventricular outflow tract obstruction to a systolic gradient of 80 mm Hg. Using the same operative procedure associated with a fibromyectomy, we completely

Fig. 1. Postoperative echocardiogram (parasternal longitudinal view): left AV valve augmentation at the level of the aorta-left AV valve continuity showing the polytetrafluoroethylene patch (arrow). AO, Aorta; LA, left atrium; LV, left ventricle; RV, right ventricle.