distal to the aortic crossclamp, that is, after ischemia. The different results are probably related to how fast the memantine can cross the blood-brain barrier. Had the intra-arterial infusion been performed over a 30-minute period before aortic crossclamping, which of course is impractical, the results might have been different. We have found that a 20- to 30-minute period is usually necessary for effects of pharmacologic agents (including those of adenosine and adenosine triphosphate) to significantly appear in the brain interstitial fluid after intravenous administration (bolus or infusion).

Tadaomi-Alfonso Miyamoto, MD
Research Department
Kokura Memorial Hospital
Kitakyushu 803-8555, Japan
Koho-J. Miyamoto, MD
II Department of Physiology
University of the Ryukyus
School of Medicine
Okinawa, Japan

REFERENCES

1. Robinson MB, Coyle JT. Glutamate and related acidic excitatory neurotransmitters: from basic science to clinical application. *FASEB J* 1987;1:446-55.
2. Buchan A, Pulsinelli WA. Hypothermia but no NMDA antagonist MK-801 attenuates neuronal damage in gerbils subjected to transient global ischemia. *J Neurosci* 1990;10:311-6.
3. Miyamoto TA, Miyamoto KJ, Ohno N. Objective assessment of CNS function within 6 hours of spinal cord ischemia in rabbits. *J Anesth* 1998;12:189-94.
4. Weiss J, Goldberg MP, Choi DW. Ketamine protects cultured neocortical neurons from hypoxic injury. *Brain Res* 1986;380:186-90.

12/8/100748

Reply to the Editor:

My colleagues and I thank Miyamoto and Miyamoto for their interest and comments on our recent article regarding the *N*-methyl-D-aspartate (NMDA) blocker memantine.¹

As we are aware of the protective effect of hypothermia on the release of excitatory amino acids, we have tried to keep the rectal temperature in a normothermic range assuming that the rectal temperature will reflect more accurate spinal cord temperature than esophageal temperature. However, we will additionally monitor esophageal temperature in future experiments.

With respect to our anesthesia protocol, a comparable experiment investigating the influence of anesthetics on neurologic recovery showed that ketamine treatment reduced acute paraplegia but increased delayed paraplegia.² In our study, the control group did not show a protective effect of ketamine compared with the groups treated with memantine. Volatile anesthetics and nitrous oxide are suppressive to myogenic motor-evoked responses,³ but ketamine is recommended as a suitable supplement to opioid/nitrous oxide anesthesia in men.⁴ In our experiments we used ketamine to keep the experimental conditions constant and to avoid false positive depression of motor-evoked potentials due to the influence of an anesthetic.

In regard to the different applications of memantine, we assumed that intra-arterial infusion of memantine into the clamped aortic segment would result in better protection than administering the drug systematically. We were surprised by the fact that the systemic group did better than the intra-aortic group. We concur that this phenomenon was due to the slow crossing of memantine through the blood-brain barrier.

Marek Ehrlich, MD
Department of Cardiothoracic Surgery
Mount Sinai Medical Center
One Gustave Levy Place
Box 1028
New York City, NY 10029

REFERENCES

1. Ehrlich M, Knolle E, Ciofica R, Boeck P, Turkof E, Konetschny R, et al. Memantine for prevention of spinal cord injury in a rabbit model. *J Thorac Cardiovasc Surg* 1999;117:285-91.
2. Naslund T, Hollier LH, Money SR, Facundus EC, Skenderis BS. Protecting the ischemic spinal cord: the influence of anesthetics and hypothermia. *Ann Surg* 1992;215:409-16.
3. Zentner J, Thees C, Pechstein U, Scheufler KM, Wirkler J, Nadstawek J. Influence of nitrous oxide on motor-evoked potentials. *Spine* 1997;22:1002-6.
4. Kalkman CJ, Drummond JC, Patel PM, Takanobu S, Randall MC. Effects of droperidol, pentobarbital, and ketamine on myogenic transcranial magnetic motor-evoked responses in humans. *Neurosurgery* 1994;35:1066-71.

12/8/100747

Possible mechanism of calcification of the BioCor stentless mitral valve treated with the No-React formula

To the Editor:

I read with great interest the brief communication by Milo and associates regarding the possible mechanism of calcification of the BioCor stentless mitral valve (BioCor, Belohorizonte, Brazil) treated with the No-React anticalcification formula (Shelhigh Inc, Millburn, NJ) in 2 adolescents.

These 2 children were 13 years old at time of implantation. Both had mitral and aortic endocarditis. Both needed double valve replacement. I assume anticoagulation was not an option. Milo and associates elected to implant the BioCor stentless aortic and mitral valves treated with No-React anticalcification formula in both children. The BioCor mitral valve was a newly designed porcine mitral valve specially fitted to be implanted in the mitral position. Milo and colleagues can be congratulated for the proficient implantations of the double valves in both children.

Mechanism of calcification. Once a valve is inflamed or infected, calcification will occur in a few months or a few years. Furthermore, nonmoving cusps tend to calcify early. The BioCor stentless mitral valve is still an experimental valve design (Fig 1). The major drawback is the length limitations of the chordae. If the patient has a large heart, the chordae will be too short. This mismatch of chordal length can cause extra tension and rupture of the chordae. Further-