LETTERS TO THE EDITOR

Actuarial and Kaplan-Meier survival analysis: There is a difference

To the Editor:

While attending the Seventy-ninth Annual Meeting of The American Association for Thoracic Surgery in New Orleans, I was dismayed to see the almost universal misuse of the term “actuarial survival.”

There are two general types of analysis for survival information: actuarial and Kaplan-Meier. An actuarial analysis should be performed when the actual date of a survival event is unknown. The known information is that the event occurred between time t_n and time t_{n+1} . Actuarial analysis is carried out at specific time intervals (6 months, 1 year), and the resulting graph will step only at those intervals. As the actual failure time is only approximated by the end point, the convention is to attribute a survival time of the fully completed intervals plus half the time of the interval during which the event occurred. Examples of events needing actuarial analysis include population-based death rates and disease-free survival.

Kaplan-Meier analysis is used when the actual date of the end point is known. End points not reached are treated as censored at the date of last follow-up for the analysis. Kaplan-Meier analysis is undertaken at each survival event, death, or censoring, and the graphs will step at each failure time and may or may not be drawn to show the location of censored observations. Examples of an appropriate event for Kaplan-Meier analysis would be postoperative survival when the date of deaths is known.

Most of the misuse of the term “actuarial survival” came during presentation of data with well-known end points and graphs that clearly reflected a Kaplan-Meier analysis. To mislabel the precise survival estimates with the term “actuarial survival” suggests a limited understanding of survival analysis.

David W. Wormuth, MD, MPH
Division of Cardiothoracic Surgery
Strong Memorial Hospital
601 Elmwood Ave
Box Surg
Rochester, NY 14642
12/8/102480

Reply to the Editor:

We are indebted to Dr Wormuth for drawing attention to inaccurate terminology related to so-called survival analysis that has crept into our Association’s annual meeting (and into submitted manuscripts). Communication is the essence of such meetings and of the *Journal*; yet communication controversies continue even within cardiothoracic surgery in the area of medical terminology. These pale, however, in comparison with those in the arena of survival analysis. When presenters and authors deem it necessary to select an adjective

to qualify their estimates of time-related events, they find no universally accepted term! Even the general name for the analyses performed is not agreed on, in part because the methodology is widely applicable: survival analysis, survival modeling, survival data analysis, actuarial analysis, life table, life-history analysis, life data analysis, life-testing, failure-time data, event-history analysis, and censored data analysis (I may have missed some). We see survival analysis in the narrow context of death and other time-related morbid events in cardiothoracic surgery; however, historically, it has roots in other disciplines that continue to contribute to its development, including demography, annuities and insurance, and life-history of machinery in industry. These are further generalized to competing risks, multiple decrement, Markov process, and other names, not to mention the host of names associated with biomathematical models! The terms have come from different, often independent, historical roots, but they all relate to the general mathematical and statistical theory of counting processes (martingales).

History. The word *actuarial* comes from the Latin *actuar-ius*, secretary of accounts. The most notable actuary was the Praetorian Prefect Domitius Ulpianus, who produced a table of annuity values early in the 3rd century AD.¹ This table continued to be used in Europe through the 18th century and even into the early 19th. With the emergence of both solid population data and the science of probability, modern so-called life tables were produced by Edmund Halley² (of comet fame) in 1693. He was motivated, as was the actuary Ulpianus, by economics as related to human survival (annuities, life insurance). Workers in this combined area of demography and economics came to be called *actuaries* in the late 18th century. Importantly for this discussion, the methodology of the actuary varied widely. In the 19th century the actuary of the Alliance of London, Benjamin Gompertz,³ developed physiologically based mathematical models of the dynamic human processes of birth and death to characterize survival. This model-based, completely parametric (equations with constants estimated from data) methodology was substantially different from the simple empiric counting methodology of Halley.

In the 300 years since Halley, a multitude of methods has been developed, and often reinvented, in actuarial science, demography, statistics, industry, and medical science. They all have the common goal of estimating the distribution of the intervals between a designated time zero and the occurrence of an event. In modern times, they also imply a suite of methodology applicable to incomplete data. That is, they permit estimates of at least portions of the distribution to be made when, for many subjects, the time of the event’s occurrence is only known to be beyond the last time of observation (so-called right censoring, one of several types of incomplete data).

Estimators versus adjectives. With this background I come to the crux of Dr Wormuth’s concerns. When authors

speak or write of actuarial survival, are they communicating inaccurately the method or formula applied to data to obtain survival estimates (the estimator), or are they simply using a historically rooted generic adjective to identify that the estimates are based on time-to-event data? My personal opinion is that they are using an adjective to identify a general type of data and its analysis rather than identifying a specific estimator. Dr Wormuth believes that they are specifying the estimator. Regardless, we routinely advise authors against use of any adjective in front of the word "survival" (and against the use of the further qualifier "rate") because it is unnecessary. They can simply state "Survival was...." The "Methods" section of the manuscript should identify the estimator used to obtain the survival estimates.

The Kaplan-Meier estimator. As Dr Wormuth states, these days the most commonly used estimator for survival and other time-related morbid events in medicine is the product-limit estimator described by Kaplan and Meier⁴ and called the Kaplan-Meier estimator. It represents the unlikely synthesis of its two authors, one working with life-times of vacuum tubes in the repeaters in telephone cables buried in the ocean and the other working with medical follow-up studies. They were forced to generate a joint paper by the editor of the *Journal of the American Statistical Association*, who recognized that their individual submissions on the subject of "failures" in the face of incomplete information were really about the same subject.

Its advantage is a firm basis in probability and statistical theory, widespread availability in commercial statistical computer programs, and its visual appeal of presenting the most atomic details of the distribution of times to events. We encourage authors to use fully the estimates obtained by application of this methodology in their graphs, for they provide more visual information than does unnecessary coarse collapsing of the estimates into interval values (as would be the product of the so-called actuarial estimator).

I would also quibble with the statement that "Kaplan-Meier analysis is undertaken at each survival event, death, or censoring." In fact, the Kaplan-Meier estimator provides survival estimates only at uncensored times (death) and not at censored times. Chin Long Chiang⁵ described a method that accounts explicitly for the times of censoring, but for some reason this more complicated method (requiring more than counting) has not seen widespread implementation.

Finally, whether or not to connect Kaplan-Meier survival estimates at all, to use a zero order interpolation (straight line) as suggested by Dr Wormuth, or first-order or higher interpolation (linear interpolation, spline interpolation) is not a settled issue. Dr Yang-Ming Zhu in my group demonstrated that both maximum entropy estimation and minimum norm estimation favor at least some form of interpolation greater than zero (step function) when using a nonparametric estimator such as that of Kaplan and Meier.

The actuarial estimator. What seems generally agreed on is that, in contrast to the Kaplan-Meier estimator, the actuarial estimator generates survivorship (freedom from events)

estimates within time intervals, rather than at the time of each event. However, the limit as the interval approaches zero is the Kaplan-Meier estimator, as was described by Böhmer in 1912, long before the seminal paper by Edward Kaplan and Paul Meier.

If I may be allowed to correct Dr Wormuth, the actuarial estimator was, and is, often used in settings in which the exact times of the events are known. For large populations, grouping of times to death was convenient and saved computational time. Certainly the introduction of the actuarial estimator to the medical mainstream by Berkson and Gage⁶ at the Mayo Clinic was not motivated by having interval data, but by not having readily available a better method! They called the method the actuarial method because they had obtained it from the 1922 writings of Murphy and Papps for the Actuarial Society of America. I agree with Dr Wormuth that, in most medical settings today, this interval method is considered obsolete and more exact methods (such as the Kaplan-Meier or Nelson-Aalen⁷ estimators) should be used. Further, the word *actuarial* carries with it the historical connotation of economic inferences from the survival data, which is generally not the case in medical follow-up reporting.

Today, the exact time of an event may not be known, only that it occurred after some point in time and before a later time point. This is known as interval-censored data. All modern life-table methods (for lack of a generic term for them), including the method of Kaplan and Meier, can be modified to accommodate a mixture of data with exact times of events known and interval-censored data. Parametric methods allow mixtures of data representing additional censoring types.⁸

Summary. Accuracy and clarity in communication are important. However, this lengthy response to the issues raised by Dr Wormuth indicates that communication within the subject of so-called survival analysis is not simple! There is lack of an accepted generic adjective for analyses of time-related events. There is historical baggage that suggests that in most medical contexts "actuarial" has implications that are not intended, although its introduction by Berkson at the Mayo Clinic is compelling. One must distinguish between the type of data at hand, the estimator applied to that data, and the estimates thereby generated.

I dare say, in all these regards and including Dr Wormuth and me, that we all have a "limited understanding of survival analysis" (for lack of a better phrase with which to characterize it!).

Eugene H. Blackstone, MD
Associate Editor

REFERENCES

1. Hacking I. The emergence of probability: a philosophical study of early ideas about probability, induction and statistical inference. Cambridge: Cambridge University Press; 1975. p. 5, 111.
2. Halley E. An estimate of the degrees of mortality of mankind, drawn from curious tables of the births and funerals at the city of Breslau; with an attempt to ascertain the price of annuities upon lives. *Phil Trans R Soc* 1693;17:596-610;654-6.
3. Gompertz B. On the nature of the function expressive of the law

- of human mortality, and on a new method of determining the value of life contingencies. *Phil Trans R Soc* 1825;27:513-85.
4. Kaplan EL, Meier P. Nonparametric estimation from incomplete observations. *J Am Stat Assoc* 1958;53:457-81.
 5. Chiang CL. The life table and its applications. Malabar: Robert E. Krieger Publishing Company; 1984. p. 221-43.
 6. Berkson J, Gage R. Calculation of survival rates for cancer. *Mayo Clin Proc* 1950;25:270-86.
 7. Nelson W. Applied life data analysis. New York: John Wiley; 1982.
 8. Hazelrig JB, Turner ME Jr, Blackstone EH. Parametric survival analysis combining longitudinal and cross-sectional-censored and interval-censored data with concomitant information. *Biometrics* 1982;38:1-15.

12/8/102963

Cardioplegic solutions: Unproved herbal approach versus tested scientific study

To the Editor:

The two criteria for cardiac surgical success are a technically adequate procedure and safe myocardial protection. Experienced surgeons can readily introduce satisfactory technical changes that visibly work. For example, a coronary anastomosis can be made with interrupted or running sutures, by passing a needle either from the outside to inside, or inside to outside, and either one surgeon doing all, or two surgeons, each doing half, to maintain an ongoing forehead direct suturing method. The barometer is that the anastomosis is hemostatic, is patent, and remains open at follow-up examination. This end point is clear and documentable.

These specific limitations with cardioplegia may not be clear in the operating room without this knowledge. For example, an untested solution may produce a minor decrease in protection, resulting in raised left ventricular end-diastolic pressure (or left atrial pressure increases from 8 mm Hg to 16 mm Hg) with normal cardiac output in normal hearts. This depression of the Starling function curve is not clear from baseline resting measurements. However, cardiac output may not increase sufficiently during anemia or fever. Of course, these changes can occur also with tested solutions with any form of myocardial protection, especially in damaged hearts. Recognition of changes by a solution alone allows more proper selection of crystalloid or blood cardioplegic constituents.

The surgeon is not qualified to make arbitrary changes in solutions to meet a whim. Suspicion of a useful change is only a reason for pharmacologic testing. This was stated clearly by Claude Bernard: "Clinical anecdotes should be the seeds of subsequent investigative studies, and not the basis of future decision making."

The calcium concentration (Ca^{++}) is lowered in the cardioplegic solution and reperfusate, because intracellular Ca^{++} accumulation follows ischemic and reperfusion injury. However, Ca^{++} is not lowered too severely ($<50 \mu\text{m}$) to avoid sarcolemmal damage that is caused by the calcium paradox.

This deleterious change was seen at the outset of clinical cardioplegia with the Melrose solution in 1955. Careful

experimental testing (as described above) was not done. We now know that Ca^{++} was lowered severely when citrate-phosphatedextrose (CPD) solution (which chelates calcium) was added in high concentrations. Cardioplegia was initially abandoned and then subsequently restored when Bretschneider in 1964 and Gay and Ebert in 1971 implemented experimental studies and showed clear benefits. Today, our current blood cardioplegic solution uses safer potassium and citrate components than were used in the original Melrose solution. The quantity of blood/cardioplegia mixture is recorded, and Ca^{++} is lowered to 0.5 to 0.6 $\mu\text{mol/L}$ with a 4:1 mixture and more severely, to 0.2 to 0.3 $\mu\text{mol/L}$, in solutions for energy-depleted hearts or those with acute myocardial infarction.

Similar testing with crystalloid Bretschneider solution or St Thomas' Hospital solution exposed the potential advantages and disadvantages of these approaches. Consequently, the literature contains comparisons that allow surgeons to make more knowledgeable selections of a desired blood or crystalloid solution.

I will relate specific issues with blood cardioplegia to show that inaccurate choices can be problematic. Several years ago, in children, we saw immediately after aortic unclamping, transient atrioventricular dissociation or ventricular fibrillation. Normal rhythm was readily restored by pacing or defibrillation. The spontaneous beating empty state seen in adults was not routine. We found that our hemodilution primes diluted Ca^{++} to 0.6 to 0.8 μm in the blood component of our cardioplegic solution. This iatrogenic hypocalcemia developed after extracorporeal circulation was begun and before the cardioplegic mixture was made. Clearly a further lowering of Ca^{++} occurred with any cardioplegic solution containing a fixed CPD concentration.

The lowering depended on the pump prime, with its known content of Ringer's lactate or electrolyte solution (Plasma-Lyte). A supplemental plasma component will lower Ca^{++} further. We worked out the formulation of the amount of calcium to be added to the crystalloid or crystalloid/plasma pump prime to restore Ca^{++} to normal limits. These transient arrhythmias were avoided by restoring normal Ca^{++} in the pump prime.¹

A more serious problem occurred at another institution. The surgeon did not dilute our 4:1 cardioplegic solution, but administered the entire cardioplegic component (containing high potassium and CPD) directly into the heart. A problem similar to that with the Melrose solution occurred, and a patient died. I² reported this hazard in a letter partially titled, "Cardioplegic Solutions Are Not Equal."

St Thomas' Hospital solution (ie, a crystalloid solution) was described in that report to show that this solution can be delivered directly into the heart without blood. Our blood cardioplegic components were also presented to show that the crystalloid components of blood cardioplegia would differ depending on the ratio of 8:1, 4:1, 1:1, or other values. Clearly, a reverse problem occurs by mixing crystalloid St Thomas' Hospital solution with blood, unless the electrolyte concentrations are increased.