LETTERS TO THE EDITOR

Off-pump internal thoracic artery–left anterior descending coronary artery grafting via small anterior thoracotomy: When and compared to what?

To the Editor:

We read with interest the article by Calafiore and associates on the midterm results after minimally invasive coronary surgery (J Thorac Cardiovasc Surg 1998;115:763-71).

The authors deserve credit for having popularized grafting of the left internal thoracic artery (ITA) to the left anterior descending coronary artery (LAD). There is some concern, however, about patient selection and patient mix in the study population, on which the analyses in the paper are based. It is uncommon to find an indication for single LAD grafting in patients with multiple-vessel disease. Indeed, on page 764 the authors state that patients who are eligible for such an operation form a "very select" group of all patients having multiple-vessel disease. It is therefore surprising that the authors managed to find 220 patients of this rare category selected for their study, an even larger group than the single-LAD group (214 patients). The total number of patients who underwent coronary artery bypass grafting in that period was not mentioned. Also, there may have been some special referrals. More important, in the presentation of the results the two groups are largely combined.

This makes generalization of the results questionable. The extraordinary nature of this large group of patients with multiple-vessel disease, treated with single-LAD grafting, is illustrated by the apparent absence of a material difference in functional outcome. It should be noted, however, that this pertains to midterm outcome only. Differences in outcome may well become visible on long-term follow-up.

The authors should consider presenting their data strictly separately for single- and multiple-vessel groups. The multiple-vessel group in the present study should be regarded with caution, because this necessarily is a highly atypical group of patients. If not, the findings challenge the basic concept of surgical myocardial revascularization, that is, complete revascularization. Still, the long-term follow-up of this atypical group is of clear interest. In its present form the results of the report are difficult to interpret and may lead to confusion.

Erik W. L. Jansen, MD, PhD

Rick E. Grobbee, MD, PhD

Department of Cardiopulmonary Surgery

Heart Lung Institute

Utrecht University Hospital (Rm E03.406)

PO Box 85500

Utrecht, 3508 GA, The Netherlands

12/8/97383

Reply to the Editor:

The relatively high number of patients with multiple-vessel disease in the population that underwent single LAD grafting described in our recent article (J Thorac Cardiovasc Surg 1998;115:763-71) disturbed more than one surgeon. The idea

Fig 1. Coronary angiogram showing a severely diseased LAD. The huge marginal branch is free of significant lesions, but the distal circumflex artery shows a severe stenosis followed by a small terminal branch.

to leave a diseased coronary vessel ungrafted seems to be a strategic mistake.

However, the inclusion of this "very select group" of patients depends on the definition of "diseased vessel" and of "complete revascularization."

My definition of "diseased vessel" is the following: A vessel is considered "diseased" if it is the main vessel of the territory or if a side branch of it is occluded or if it has a lesion equal to or greater than 70%. If a circumflex artery or a right coronary artery has a short and small posterolateral branch with such lesions, the circumflex artery or the right coronary artery has to be defined as "diseased." In Fig 1, a coronary angiogram is shown in which the LAD has severe disease; the huge marginal branch is free of significant lesions, but the distal circumflex artery shows a severe stenosis followed by a small terminal branch. This patient has 2-vessel disease, but I would never graft this small branch which, in my opinion, has no influence on the patient's future. I certainly would never use cardiopulmonary bypass for such a purpose.

The concept of "complete revascularization" needs a more thorough analysis. If with this term we mean that any lesion that causes ischemia to the patient should be grafted, I fully agree. If this definition also includes the necessity to graft major and important coronary vessels supplying a huge territory and with a significant ($\geq 70\%$) lesion, even if there is no clearly demonstrated ischemia, when concomitant grafting to other surely ischemic territories is performed, I fully agree.

But if with this term we state that any stenotic vessel has to be grafted, regardless of its importance and size, without any demonstration of ischemia in the territory, this definition of "complete revascularization" has to be forgotten. Today we have at our disposition many tools to properly investigate the importance of a stenosis, and complete revascularization (interventional, surgical, or combined) of all ischemic territories is always our goal. However, the timing can change and the strategy can differ from patient to patient.

In our article I thought that the real problem was focused not on the surgical indication (single LAD disease or multiple-vessel disease), which can be debatable, but on the future of the graft. (Is this approach able to guarantee satisfying permeability of the graft? Can this strategy be considered safe and reproducible?) For this reason I mixed all my patients, but the patients with "multiple-vessel disease" in this group, with only a few exceptions (hybrid procedures), are not the same patients we operate on every day in our daily practice.

Antonio Maria Calafiore
Department of Cardiac Surgery
San Camillo de' Lellis Hospital
Via Forlanini 50
Chieti 66100, Italy
12/8/97384

The use of the calcium antagonist nicardipine in arterial coronary bypass surgery

To the Editor:

In the December 1998 issue of the Journal, Acar,¹ Possati,² and their associates independently provided important data on the 5-year patency rate of the radial artery graft.

Both groups acknowledge the impressive spastic characteristics of this arterial conduit and advocate perioperative release of spasm and routine administration of the calcium antagonist diltiazem to the patients for 1 year or more. Interruption of this treatment after 1 year or later during follow-up did not seem to adversely influence either clinical outcome or 5-year patency. This observation is related to the apparent loss of reactivity of the radial graft over time.

The use of antispasmodic drugs for patients receiving arterial grafts is a common practice, although so far no study has demonstrated its value on the surgical outcome, keeping in mind that "spasm" may be confounded by other factors such as surgical technique, familiarity with arterial grafts, and perioperative conduit preparation. Since 1994, we have been using nicardipine hydrochloride, the first intravenously administered dihydropyridine calcium channel antagonist, for sequential or bilateral internal thoracic artery grafting. The patients are started perioperatively on intravenous nicardipine hydrochloride (0.25 µg/kg per minute) after a 1 mg intravenous bolus. The drug is titrated according to the systemic vascular resistance and discontinued on the second postoperative day. For patients receiving a radial artery graft, we added to this protocol gentle hydrostatic dilatation of the conduit with 1% papaverine and oral administration of nicardipine 20 to 30 mg three times a day after discontinuation of intra-

venous perfusion on day 2. The first 50 patients undergoing myocardial revascularization with a radial graft using that protocol were observed. A total of 150 anastomoses were performed with a mean of 3.0 anastomoses per patient, and 108 arterial grafts were used for the completion of 111 coronary distal anastomoses. Fifty radial arteries were used for 52 distal anastomoses, and 36 venous grafts were used for 39 distal anastomoses. The radial artery was placed on the obtuse marginal branch in 58% of cases, diagonal branches in 13%, and the right coronary artery in 29%. Proximal anastomoses were done directly on the aorta in 80% of cases.

The operative mortality was 4% because of 2 cases of fulminant pulmonary sepsis. There was no evidence of perioperative myocardial infarction or arterial graft hypoperfusion syndrome. Mean creatine kinase MB level at 18 hours was 36 µg/L. Neither ischemic anomalies of the electrocardiogram nor wall motion abnormalities on discharge transthoracic echocardiography were detected. Angiography performed in the last 20 patients showed a 98% (51/52) permeability rate for all grafts. Nineteen of 20 radial grafts were patent. A moderate spasm (40%) developed in the middle part of the conduit in one radial artery.

Our experience with the radial artery is part of a larger experience with the perioperative use of nicardipine in more than 550 cases of bilateral or sequential internal thoracic artery grafting. This protocol has virtually eliminated internal thoracic artery spasm or hypoperfusion in our practice. Nicardipine has several potential advantages over diltiazem. Nicardipine is a more potent and selective arteriolar vasodilatory agent, which also has the capability to inhibit endothelin-induced vasoconstriction.³ This drug has neither chronotropic, dromotropic, nor inotropic negative effects. Its short action duration makes it convenient for perioperative management, and it is well tolerated in association with β blockers in the routine management of patients with coronary artery disease. Nicardipine also has a documented cardioprotective effect on ischemic myocardium, and some data suggest cardioprotective effects during cardioplegic cardiac arrest.⁴ We therefore suggest that the time has come to assess the real need for anti-spasmodic drugs in arterial coronary bypass grafting and to determine the most adequate pharmacologic management for patients receiving arterial grafts such as the radial artery.

M. A. Radermecker, MD^a

T. Grenade, MD^a

R. Larbuisson, MD^b

R. Limet, MD^a

Department of Cardiovascular Surgery^a

Department of Anesthesiology^b

CHU du Sart-Tilman

4000 Liege, Belgium

REFERENCES

1. Acar C, Ramsheji A, Pagny JY, Jebara V, Barrier P, Fabiani JN, et al. The radial artery for coronary artery bypass grafting: clinical and angiographic results at five years. *J Thorac Cardiovasc Surg* 1998;116:981-9.
2. Possati G, Gaudino M, Alessandrini F, Luciani N, Glietta F, Trani C, et al. Midterm clinical and angiographic results of radial artery