pletely 6 hours after reperfusion.² In their study, 30% of animals subjected to 15 minutes of ischemia recovered function 8 hours after reperfusion, and 1 rabbit that had not recovered 24 hours after reperfusion did recover 2 days later. For 15 minutes of ischemia to result in 30% recovering acutely and 40% of animals recovering normal function when evaluated 2 days later, some sort of a protection (temperature or anesthetic factors, or a combination) must have taken place. Their anesthesia included ketamine (50 mg/kg), and ketamine is known to be an *N*-methyl-D-aspartate receptor antagonist, which in general promotes and facilitates hypothermia as part of its protective effects.^{1,3,4} Since data obtained from experiments in which ketamine is used could not be extrapolated to the clinical situation, in which ketamine is not used, and because knowing the ischemic times when apoptosis starts to occur would be of clinical significance, the authors are encouraged to repeat their experiments without ketamine. We would suggest that they use volatile agents at concentrations that are not protective by themselves, rather than intravenous or intramuscular agents, and perhaps shorter ischemic times (6 and 7 minutes). Our prediction is that 7 minutes of ischemia at 38.5°C should result in a good number of animals showing apoptosis if seen 2 days after the ischemic episode. Also, although unlikely (since we believe apoptosis is a mechanism to get rid of injured elements), it would not be surprising if some of the animals with 6 minutes (so far found to recover completely within 6 hours of reperfusion) may prove to have apoptotic late neuronal death despite not having evidence of functional impairment when seen 6 hours after reperfusion.

Tadaomi-A. Miyamoto, MD
Research Department
Kokura Memorial Hospital
Kitakyushu, Japan
Koho-J. Miyamoto, MD, PhD
Assistant Professor
II Department of Physiology
University of the Ryukyus
School of Medicine
Okinawa, Japan

REFERENCES

1. Buchan A, Pulsinelli WA. Hypothermia but not NMDA antagonist MK-801 attenuates neuronal damage in gerbils subjected to transient global ischemia. *J Neurosci* 1990;10:311-6.
2. Miyamoto TA, Miyamoto KJ, Ohno N. Objective assessment of CNS function within 6 hours of spinal cord ischemia in rabbits. *J Anesth* 1998;12:189-94.
3. Weiss J, Goldberg MP, Choi DW. Ketamine protects cultured neocortical neurons from hypoxic injury. *Brain Res* 1986;380:186-90.
4. Robinson MB, Coyle JT. Glutamate and related acidic excitatory neurotransmitters: from basic science to clinical application. *FASEB J* 1987;1:446-55.

12/8/96848

Reply to the Editor:

In our model, systemic proximal aortic pressure revealed only a trivial and transient change, not a significant change.

Hence we believe that the change in proximal pressure has no influence on the spinal cord injury.

In this model spinal temperature was 37°C. Miyamoto and Miyamoto cited a reference that spinal cord temperature was more closely correlated with esophageal temperature. However, the correlation was not in the lumbar region but in the thoracic region, since the esophagus and lumbar region are at different levels. We believe esophageal temperature does not precisely reflect the lumbar spinal cord temperature. Therefore, we measured the temperature of the rectum, which is more proximal to the spinal cord in the lumbar region.

The models described by Miyamoto and Miyamoto are quite invasive. Previous reports describing spinal cord ischemia were made using laparotomy^{1,2}; however, the aorta was occluded from 4 hours to 2 days after closure of the abdominal wall (the authors did not administer ketamine during the time the aorta was occluded). Therefore, their spinal ischemia model may not recover after 8 or 9 minutes of ischemia.

The word "apoptosis" describes histologic and biologic features.³⁻⁶ However, in their letter, Miyamoto and Miyamoto have not described histologic and biologic findings in spinal cord ischemia.

Masahiro Sakurai, MD
Takeshi Hayashi, MD
Tohoku University School of Medicine
Sendai, Japan
Koji Abe, MD, PhD
Okayama University School of Medicine
Okayama, Japan
Mitsuaki Sadahiro, MD, PhD
Koichi Tabayashi, MD, PhD
Tohoku University School of Medicine
Sendai, Japan

REFERENCES

1. Moore WM, Hollier LY. The influence of severity of spinal cord ischemia in the etiology of delayed-onset paraplegia. *Ann Surg* 1991;213:427-32.
2. Zivin JA, DeGirolami U, Hurwitz EL. Spectrum of neurological deficits in experimental CNS ischemia: a quantitative study. *Arch Neurol* 1982;39:408-12.
3. Nitatori T, Sato N, Waguri S, Karasawa Y, Araki H, Shibana K, et al. Delayed neuronal death in the CA1 pyramidal cell layer of the gerbil hippocampus following transient ischemia is apoptosis. *J Neurosci* 1995;15:1001-11.
4. Héron A, Pollard H, Dessi F, Moreau J, Lasbennes F, Ben-Ari Y, et al. Regional variability in DNA fragmentation after global ischemia evidenced by combined histological and gel electrophoresis observations in the rat brain. *J Neurochem* 1993;61:1973-6.
5. Li Y, Chopp M, Yao F, Zaloga C. Temporal profile of in situ DNA fragmentation after transient middle cerebral artery occlusion in the rat. *J Cereb Blood Flow Metab* 1995;15:389-97.
6. Arends MJ, Wyllie AH. Apoptosis: mechanisms and roles in pathology. *Int Rev Exp Pathol* 1991;32:223-54.

12/8/96847